

**■ BASF**

We create chemistry

# Ornamental Plant Production Guide

Your Guide to Healthy, Quality Plants



# contents

- 1 **Innovate. Collaborate. Grow.**
- 2 **Ornamental Plant Production Product Use Site Overview**
- 7 **Disease Management**
  - 7 Fungicide Use Sites
  - 8 When to Use BASF Fungicides In The Ornamentals Production Cycle
  - 9 The Plant Health Story Behind **Intrinsic®** Brand Fungicides
  - 10 Disease Management Products
  - 17 Fungicide Resistance Management Guide
- 25 **Mite and Insect Management**
  - 25 Insecticide Use Sites
  - 26 Mite and Insect Management Products
  - 36 Insecticide Resistance Management Guide
- 43 **Weed Management**
  - 43 Herbicide Use Sites
  - 44 Weeds Controlled
  - 45 Weed Management Products
  - 53 Herbicide Resistance Management Guide
- 59 **Total Release Fungicides and Insecticides**
  - 59 How to Use Total Release Products
  - 61 Total Release Products Chart
- 67 **Specialty Greenhouse Products**
  - 67 Specialty Greenhouse Product Use Sites
  - 68 Iron Chelates
  - 70 Disinfectant and Algicide
  - 71 Insect Monitoring and Trapping Cards
  - 72 Spray Pattern Indicator
  - 73 Tank Cleaners
  - 74 Wasp and Hornet Insecticide

## **Innovation that solves your toughest challenges**

*BASF is dedicated to investing in research that produces quality solutions. We develop innovations that help solve your most pressing challenges, backed by research and supported by an experienced sales and technical team. We create chemistry that leads to healthy, quality plants, greater efficiency and more sustainable solutions for greenhouse and nursery growers.*



## Innovate. Collaborate. Grow.

Greenhouse and nursery growers are innovators, always looking to improve operational efficiency, learn about the latest solutions, and implement advancements. BASF is at your side, listening to your challenges and working with you to deliver information and solutions as innovative as you are. Together, we will continue to find common ground and collaborate. Together, we will grow.

### How to Use This Guide

This comprehensive reference guide includes all the BASF products and use guidelines needed to ensure healthy, high-quality plant performance. Divided into five sections, it organizes BASF solutions according to areas of plant protection: Disease Management; Mite and Insect Management; and Weed Management, with additional sections dedicated to Total Release Fungicides and Insecticides, and Specialty Greenhouse Products.

Within each section, you will find reference resources such as Use Site Recommendations, Spectrum of Control Overviews, Resistance Management Guides and summarized product information, complete with QR (Quick Response) code links to CDMS, the authoritative source for label information. The product label of any companies' products remains the definitive source for how a product is used and registered. Please read and follow specific label directions and restrictions.

Scan the QR code below, or visit [betterplants.basf.us](https://betterplants.basf.us) for more detailed technical information on every BASF product listed in this guide.



# Ornamental Plant Production Product Use Site Overview

## Fungicide Use Sites

Product	Greenhouse	Container Nursery	Field Production	Conifer Nursery Plantation	Retail Greenhouse / Nursery	Interiorscape	Landscape and Recreational Areas	Vegetable Transplants and Production
<b>Intrinsic® Brand Fungicides</b>								
<b>Pageant® Intrinsic</b> Brand Fungicide	■	■	■	■	■	■	■	■
<b>Empress® Intrinsic</b> Brand Fungicide	■	■	■	■	■	■		
<b>Orkestra® Intrinsic</b> Brand Fungicide	■	■	■	■	■	■	■	
<b>Specialty Fungicides</b>								
<b>Trinity®</b> Fungicide	■	■	■	■	■	■	■	
<b>Orvego®</b> Fungicide	■	■	■	■	■	■		
<b>Stature SC®</b> Fungicide	■	■	■	■	■	■		
<b>Ultra Pure Oil®</b> Horticultural Insecticide, Miticide and Fungicide	■	■	■	■	■	■	■	■
<b>Total Release Fungicides</b>								
<b>Trinity® TR</b> Total Release Fungicide	■							
<b>Pageant® TR Intrinsic®</b> Total Release Fungicide	■*							
<b>Biological Fungicides</b>								
<b>Serifel® NG</b> Biological Fungicide	■	■	■	■	■			■

\* Pageant TR Intrinsic brand fungicide may also be used in trucks - cargo area only.

## Insecticide Use Sites

Product	Greenhouse	Container Nursery	Field Production	Conifer Nursery Plantation	Retail Greenhouse / Nursery	Interiorscape	Landscape and Recreational Areas	Vegetable Transplants and Production
<b>Specialty Insecticides</b>								
<b>Ventigra™</b> insecticide	■	■	■	■	■	■	■	■
<b>Sultan®</b> Miticide	■	■	■	■	■	■	■	
<b>Pylon®</b> Miticide/Insecticide	■							■
<b>DuraGuard® ME</b> Microencapsulated Insecticide	■	■	■				■	
<b>Siesta®</b> Fire Ant Bait		■	■	■	■		■	
<b>Amdro® Pro</b> Insect Fire Ant Bait	■	■	■	■	■		■	
<b>Ultra Pure Oil®</b> Horticultural Insecticide, Miticide and Fungicide	■	■	■	■	■	■	■	■
<b>Biological Insect Controls</b>								
<b>Nemasys®</b> Beneficial Nematodes	■	■	■	■	■	■		
<b>Millenium®</b> Beneficial Nematodes	■	■	■	■	■	■	■	
<b>Velifer®</b> Fungal Contact Insecticide	■							■
<b>Total Release Insecticides</b>								
<b>Beethoven® TR</b> Miticide/Insecticide	■							
<b>Pylon TR</b> Total Release Miticide/ Insecticide	■							
<b>1300 Orthene® TR</b> Total Release Insecticide	■							
<b>Attain® TR</b> Total Release Insecticide	■							
<b>Duraplex® TR</b> Total Release Insecticide	■							
<b>Pyrethrum® TR</b> Total Release Insecticide	■							

## Herbicide Use Sites

Product	Bareground	Conifer Nursery Plantation	Container Nursery	Field Production	Retail Nursery	Landscape and Recreational Areas	Greenhouse and Shadehouse
<b>Preemergent Herbicides</b>							
<b>FreeHand® 1.75G</b> Herbicide	■	■	■	■	■	■	
<b>Pendulum® AquaCap</b> Herbicide	■	■	■	■	■	■	
<b>Pendulum® 3.3 EC</b> Herbicide	■	■	■	■	■	■	
<b>Pendulum® 2G</b> Herbicide	■	■	■	■	■	■	
<b>Tower®</b> Herbicide	■ <sup>1</sup>	■	■	■	■	■	■
<b>Postemergent Herbicides</b>							
<b>Finale®</b> Herbicide	■	■	■	■	■	■	■
<b>Basagran® T&amp;O</b> Herbicide	■	■	■	■	■	■	
<b>Segment® II</b> Herbicide	■	■	■	■	■	■	

## Specialty Greenhouse Product Use Site Recommendations

Product	Bareground	Conifer Nursery Plantation	Container Nursery	Field Production	Retail Nursery	Greenhouse	Interior	Landscape and Recreational Areas	Turf
<b>Pest Control</b>									
<b>PT® Wasp Freeze®</b> Wasp & Hornet Insecticide	■	■	■	■	■	■	■	■	■
<b>Iron Chelates</b>									
<b>Sprint® 138</b> Iron Chelates		■	■	■	■	■	■	■	■
<b>Sprint® 330</b> Iron Chelates		■	■	■	■	■	■	■	■
<b>Disinfectant and Algicide</b>									
<b>Green-Shield® II</b> Disinfectant and Algicide						■	■ <sup>2</sup>		
<b>Insect Monitoring</b>									
<b>Sensor®</b> Insect Monitoring and Trapping Cards		■	■	■	■	■	■		
<b>Spray Pattern Indicators</b>									
<b>Turf Mark®</b> Spray Pattern Indicator	■	■	■	■		■		■	■
<b>Tank Cleaners</b>									
<b>Nutra-Sol®</b> Tank Cleaner	■	■	■	■	■	■	■	■	■
<b>Neutralize®</b> Tank Cleaner	■	■	■	■	■	■	■	■	■
<b>Other</b>									
<b>Instant Soap 'N Water®</b> Foaming Cleanser <sup>3</sup>	■	■	■	■	■	■	■	■	■

<sup>1</sup> Tower herbicide label allows for use on gravel or ground floors of open-sided lathhouses (shadehouses) or other polyhouse structures that allow polycovers to be removed on a seasonal basis.

<sup>2</sup> Consult Green-Shield II label for interior uses.

<sup>3</sup> Instant Soap 'N Water foaming cleanser is labeled for use on hands in all use sites listed.











# Disease Management

BASF provides broad-spectrum disease management solutions for every stage of the plant production cycle. **Intrinsic®** brand fungicides control the majority of ornamental diseases while delivering proven plant health benefits. The BASF disease management portfolio also includes specialist fungicides that control downy mildew, phytophthora, stem, crown and basal rot diseases.

## Fungicide Use Sites

Product	Greenhouse	Container Nursery	Field Production	Conifer Nursery Plantation	Retail Greenhouse / Nursery	Interiorscape	Landscape and Recreational Areas	Vegetable Transplants and Production
<b>Intrinsic® Brand Fungicides</b>								
<b>Pageant® Intrinsic</b> Brand Fungicide	■	■	■	■	■	■	■	■
<b>Empress® Intrinsic</b> Brand Fungicide	■	■	■	■	■	■		
<b>Orkestra® Intrinsic</b> Brand Fungicide	■	■	■	■	■	■	■	
<b>Specialty Fungicides</b>								
<b>Trinity®</b> Fungicide	■	■	■	■	■	■	■	
<b>Orvego®</b> Fungicide	■	■	■	■	■	■		
<b>Stature SC®</b> Fungicide	■	■	■	■	■	■		
<b>Ultra Pure Oil®</b> Horticultural Insecticide, Miticide and Fungicide	■	■	■	■	■	■	■	■
<b>Total Release Fungicides</b>								
<b>Trinity® TR</b> Total Release Fungicide	■							
<b>Pageant® TR Intrinsic®</b> Total Release Fungicide	■*							
<b>Biological Fungicides</b>								
<b>Serifel® NG</b> Biological Fungicide	■	■	■	■	■			■

\* Pageant TR Intrinsic brand fungicide may also be used in trucks - cargo area only.

## When to Use BASF Fungicides in the Ornamentals Production Cycle\*



### YOUNG PLANTS

#### Pageant®

Intrinsic® Brand Fungicide

Spray on unrooted cuttings, seedlings

#### Empress®

Intrinsic® Brand Fungicide

Drench on rooted plugs, seedlings, cuttings



### TRANSPLANTS

#### Empress®

Intrinsic® Brand Fungicide

Drench to prevent major root diseases



### GROWTH TO MATURITY

#### Orkestra®

Intrinsic® Brand Fungicide

Spray mid-production for fast and long-acting control of major diseases

#### Empress®

Intrinsic® Brand Fungicide

Drench to prevent major root diseases, growth to maturity



### MARKET READY

#### Pageant®

Intrinsic® Brand Fungicide

Spray for protection against foliar, stem and crown diseases through handling, shipping and in market

#### Pageant TR

Intrinsic® Brand Fungicide

Spray the aerosol in the enclosed cargo area of delivery truck for protection during transportation.

*Stature SC fungicide, Orvego fungicide and Trinity fungicide can be used from transplanting through growth to maturity.*

#### Stature® SC

Fungicide

In rotation as the first choice for treatment of downy mildew diseases

#### Orvego®

Fungicide

In rotation to prevent major root diseases of *Phytophthora* spp.

#### Trinity®

Fungicide

In rotation to prevent stem, crown and root rots and diseases

#### Ultra Pure Oil®

Horticultural Insecticide, Miticide and Fungicide

Spray for broad-spectrum control of insects, mites and fungus

\* This is not a disease resistance management program recommendation.

Consult with your BASF sales representative for a comprehensive disease management program.



## The Plant Health Story Behind Intrinsic® Brand Fungicides

At each stage of the production process, **Intrinsic** brand fungicides provide superior, broad-spectrum disease control that helps plants grow strong. This innovative line of fungicides also delivers plant health benefits, including greater tolerance to stress and increased growth efficiency for stronger, denser roots. Research shows that plants treated with **Intrinsic** brand fungicides display improved resilience to stress imparted by temperature extremes, drought, handling and transportation. The result is healthy, high-quality plants.

Start your production cycle on a strong foundation of broad spectrum disease control and resistance management. Incorporate **Pageant**® **Intrinsic** brand fungicide, **Empress**® **Intrinsic** brand fungicide and **Orchestra**® **Intrinsic** brand fungicide into your rotation to promote plant health throughout the growing cycle.





# Pageant®

## Intrinsic® Brand Fungicide

**Pageant Intrinsic** brand fungicide forms the foundation of your disease management program. It helps professional growers achieve exceptional disease control and plant health throughout propagation, production, finish operations and shipping.

### Key Features and Benefits

- Broad-spectrum, superior disease control
- Faster, stronger rooting
- Helps produce consistent, high quality plant growth

### Key Diseases Controlled

- Anthracnose
- Blossom blight
- Crown and basal rot
- Downey mildew
- Leaf spot
- *Phytophthora* aerial blight
- Powdery mildew
- Rot, blight
- Rust
- Scab
- Crown and soilborne diseases

### Use Sites

- Greenhouses
- Lathhouses and shadehouses
- Outdoor nurseries (field and container)
- Forest and conifer nurseries and plantations
- Retail nurseries
- Interiorscapes
- Golf courses
- Recreational landscapes
- Residential and commercial landscapes

Labeled for vegetable transplants for the home consumer market and vegetables grown in commercial greenhouses.

**FRAC groups:** 7 + 11

**Active ingredients:**

Boscalid  
Pyraclostrobin

**Signal word:** CAUTION

**Formulation:**

WG (Wettable granule)

**Use rates:** 4-18 fl oz per 100 gal

**PPE:** Coveralls, chemical-resistant gloves made of any waterproof material, shoes, plus socks

**REI:** 12 hours

**Packaging:** 4 x 1 lb case, 4 x 12 lb case



# Empress®

## Intrinsic® Brand Fungicide

Drench applications of **Empress Intrinsic** brand fungicide protect young, susceptible plants from root-infecting pathogens in plug, propagation and transplant stages of greenhouse and nursery production.

### Key Features and Benefits

- Outstanding control of soil-borne diseases
- Enhanced root growth
- Consistent performance
- Readily fits disease resistance management programs

### Key Diseases Controlled

Crown, root and soilborne diseases caused by:

- *Fusarium* spp.
- *Phytophthora* spp.
- *Pythium* spp.
- *Rhizoctonia* spp.

### Use Sites

- Greenhouses
- Forest and conifer nurseries
- Lathouses and shadehouses
- Interiorscapes
- Outdoor nurseries (field and container)
- Retail nurseries



**FRAC group:** 11

**Active ingredient:**

Pyraclostrobin

**Signal word:** WARNING

**Formulation:** SC (soluble concentrate)


**Use rates:** 1.0 - 6.0 fl oz per 100 gal

**PPE:** Long-sleeved shirt and long pants; chemical-resistant gloves made of any waterproof material; shoes plus socks

**REI:** 12 hours

**Packaging:** 24 oz bottle  
6 x 24 oz case





# Orkestra®

## Intrinsic® Brand Fungicide

For the nursery and greenhouse grower, **Orkestra Intrinsic** brand fungicide provides fast, long-lasting control of key diseases during the production cycle, resulting in healthier plants that exhibit the highest quality and consistency.

**FRAC groups:** 7 + 11

**Active ingredients:**

Fluxapyroxad  
Pyraclostrobin

**Signal word:** WARNING

**Formulation:** SC (soluble concentrate)

**Use rates:** 4-11 fl oz per 100 gal

**PPE:** Chemical-resistant gloves, long-sleeved shirt, long pants, shoes plus socks

**REI:** 12 hours

**Packaging:** 4 x 16 fl oz case, 4 x 80 oz

### Key Features and Benefits

- Early curative and preventive plant protection
- Translaminar and systemic activity maximizes distribution within treated plants
- Lasting protection, even under the wettest conditions
- Extends the plant health benefits of **Intrinsic** brand fungicides throughout the plant production cycle

### Key Diseases Controlled

- Anthracnose
- Crown and basal rots
- Downy mildews
- Flower and petal blights
- Leaf spots
- *Phytophthora* aerial blights
- Powdery mildews
- Rusts
- Scabs
- Stem and twig blights and dieback
- *Thielaviopsis* (suppression)
- Crown and soilborne diseases

### Use Sites

- Greenhouses
- Lathhouses and shadehouses
- Outdoor nurseries (field and container)
- Forest and conifer nurseries and plantations
- Retail nurseries
- Interiorscapes
- Golf courses
- Recreational landscapes
- Residential and commercial landscapes





# Trinity®

## Fungicide

**Trinity** fungicide provides growers and landscapers with confident control of powdery mildew, stem and crown blights, rots and other diseases in an innovative formulation that leaves no unsightly residue. **Trinity** fungicide is also available as **Trinity TR**, as shown in the Total Release section of this guide.

### Key Features and Benefits

- An effective rotation partner with **Intrinsic**® brand fungicides
- Superior plant safety compared to other DMI fungicides
- Innovative formulation; leaves no unsightly residue

### Key Diseases Controlled

- Anthracnose
- Crown and basal rots
- Leaf spots
- Powdery mildews
- Scabs
- Stem blight-diebacks

### Use Sites

- Greenhouses
- Golf courses
- Lathouses and shadehouses
- Outdoor nurseries (field and container)
- Christmas tree and conifer and/or hardwood tree nurseries
- Retail nurseries
- Residential and commercial landscapes
- Landscape beds
- Interiorscapes



**FRAC group:** 3

**Active ingredient:**

Triticonazole

**Signal word:** CAUTION

**Formulation:** SC

(Suspension Concentrate)

**Use rates:** 4 -12 fl oz per 100 gal

**PPE:** Long-sleeved shirt and long pants; chemical-resistant gloves made of waterproof material; shoes plus socks

**REI:** 12 hours

**Packaging:** 2 x 2.5 gal





# Orvego®

## Fungicide

**Orvego** fungicide meets the needs of growers who want fast knockdown to stop the spread of all major *Phytophthora* species and downy mildews in their greenhouse and nursery crops.

### Key Features and Benefits

- An effective rotation partner with **Intrinsic**® brand fungicides
- A unique mode of action for control of oomycete fungi
- Knockdown of infectious zoospore stage within seconds, stopping disease spread
- Translaminar and locally systemic activity maximizes distribution within the treated area
- Flexible foliar and drench applications for control of leaf and root diseases

### Key Diseases Controlled

- Downy mildew
- Aerial *Phytophthora*
- *Phytophthora* root, crown and stem rot

### Use Sites

- Greenhouses
- Lathhouses and shadehouses
- Hardwood and conifer nurseries  
or plantations
- Outdoor or field nurseries
- Interiorscapes

**FRAC groups:** 40 + 45

**Active ingredients:**

Ametoctradin  
Dimethomorph

**Signal word:** CAUTION

**Formulation:** Suspension

**Use rates:** 11- 14 fl oz  
per 100 gal

**PPE:** Long sleeved shirt  
and long pants, chemical-  
resistant gloves made of  
any waterproof material,  
shoes plus socks

**REI:** 12 hours

**Packaging:** 6 x 28 fl oz  
per case



# Stature® SC

Fungicide

**Stature SC** fungicide is the clear choice for the grower who wants the best preventive control over downy mildews and other oomycete diseases.

## Key Features and Benefits

- An effective rotation partner with **Intrinsic®** brand fungicides for preventive control of downy mildews and *Phytophthora* diseases
- Translaminar and locally systemic activity maximizes distribution within the treated plant for more consistent control of downy mildew
- Anti-sporulant, stops disease progression
- Flexible foliar, drench and chemigation applications for control of foliar and root diseases

## Key Diseases Controlled

- Downy mildews
- Aerial *Phytophthora*
- *Phytophthora* roots, crown and stem rots

## Use Sites

- Greenhouses
- Outdoor nurseries (field and container)
- Shadehouses and lathhouses
- Conifer seedling nurseries
- Hydroponic, rockwool or artificial substrate systems



**FRAC group:** 40

**Active ingredient:**  
Dimethomorph

**Signal word:** CAUTION

**Formulation:** SC  
(Suspension Concentrate)

**Use rates:** 3.06 to 12.25  
fl oz per 100 gal

**PPE:** Long-sleeved  
shirt and long pants;  
chemical-resistant  
gloves made of  
waterproof material;  
shoes plus socks

**REI:** 12 hours

**Packaging:** 25 fl oz,  
6 x 25 case







# Serifel® NG

## Biological Fungicide

**Serifel NG** biological fungicide controls a broad spectrum of diseases in nurseries and greenhouses by forming an invisible shield of protection on plant surfaces.

### Key Features and Benefits

- Protects against phytophthora, rhizoctonia, fusarium and pythium root pathogens early in the production cycle
- Highly effective disease control through multiple modes of action
  - Physical barrier - creates a shield on plant surfaces
  - Nutrient competition - consumes limited nutrients on plant surfaces eliminating availability to pathogens
  - Fungicidal metabolites - prevents pathogen spore germination and hyphae development
- Most concentrated biological fungicide on the market due to its pure spore formulation

### Key Pests Controlled

- Rhizoctonia spp.
- Pythium spp.
- Fusarium spp.
- Botrytis
- Powdery Mildew
  - *Podosphaera* spp.
  - *Oidiopsis* spp.
  - *Sphaerotheca* spp.
  - *Erysiphe* spp.

### Use Sites

- Greenhouse
- Field/Container Nursery

**FRAC groups:** n/a

**Active ingredients:**

Bacillus amyloliquefaciens strain MBI 600

**Signal word:** CAUTION

**Formulation:** Disbursable Powder

**Use rates:** 0.2 - 0.4 oz per 100 gallons or 1-3 # per acre

Refer to Label

**PPE:** Coveralls, chemical-resistant gloves made of any waterproof material, shoes plus socks and protective eyewear

**REI:** 12 hours

**Packaging:** 5 lb bottle

# Fungicide Resistance Management Guide

FRAC Group	Chemical Group(s)	Primary Target Disease(s)	Active Ingredient(s)	Fungicide Trade Name(s)	REI	Use Site(s)
1	<b>Thiophanate</b>	<i>Rhizoctonia</i> , <i>Thielaviopsis</i> , Fungal leaf spots	Thiophanate-methyl	AllBan®, Cleary's 3336®, OHP 6672®, T-Storm®	12	G,N,L
2	<b>Dicarboximide</b>	<i>Botrytis</i> , Powdery mildew, Fungal leaf spots, <i>Fusarium</i>	Iprodione	26GT®, Chipco® 26019	12	G,N,L
3	<b>Imidazole</b>	Scab, Powdery mildew, Fungal leaf spots, Rusts	Triflumizole	Terraguard®	12	G,N
	<b>Triazole</b>	Powdery mildews, Scab, Rusts, Blights, Fungal leaf spots	Metconazole	Tourney®	12	N,L
		Powdery mildew, Scab, Rusts, Fungal leaf spots	Myclobutanil	Eagle® 20 EW	24	G,N,L
			Propiconazole	Alamo®, Banner MAXX® II, Spectator™	24	N,L
		Powdery mildews, Scab, Rusts, Blights, Fungal leaf spots	Tebuconazole	Torque™	12	N,L
		Oak wilt, Dutch elm disease, Scab	Tebuconazole	Tebusject™	N/A	L
		Powdery mildew, Scab, Rusts, Blights, Fungal leaf spots	Triadimefon	Bayleton®	12	G,N,L
	Anthrachnose, Powdery mildew, Leaf spots, Stem and Crown rots	Triticonazole	<b>Trinity®</b> fungicide, <b>Trinity® TR</b> total release fungicide	12	G,N,L G	
4	<b>Acylalanine</b>	<i>Pythium</i> , <i>Phytophthora</i>	Mefenoxam	Mefenoxam® 2, Subdue MAXX®	0	G,N,L
5	<b>Piperadine</b>	Powdery mildew	Piperalin	Pipron®	12	G
7	<b>Phenyl-benzamide</b>	<i>Rhizoctonia</i> , Southern blight, Rusts	Flutolanil	ProStar® WG, Contrast®	12	G,N
	<b>Thiopene amides</b>	<i>Botrytis</i> , powdery mildew, <i>fusarium</i>	Isofetamid	Astun	12	GH/N

G = Greenhouse, N= Nursery, L=Landscape

1 Some products within the same chemical group may not control all pests listed, or there may be some products within the group that offer superior performance. Consult your local manufacturer's representative or your university/extension specialist if needed.

2 Specific use sites may vary by formulation and manufacturer; therefore, always read and consult product label directions.

## Fungicide Resistance Management Guide – (continued)

FRAC Group	Chemical Group(s)	Primary Target Disease(s)	Active Ingredient(s)	Fungicide Trade Name(s)	REI	Use Site(s)
11	<b>Strobilurin</b>	Fungal leaf spots, Rusts, Powdery mildew, <i>Phytophthora</i> , <i>Rhizoctonia</i>	Azoxystrobin	Heritage®	4	G,N,L
		<i>Rhizoctonia</i> , Fungal leaf spots & blights, Scab, Rusts, Anthracnose, Downy mildew, Powdery mildew, <i>Fusarium</i> , <i>Phytophthora</i> , Southern blight	Fluoxastrobin	Fame®, Disarm O	12	G,N,L
		<i>Fusarium</i> , <i>Phytophthora</i> , <i>Pythium</i> , <i>Rhizoctonia</i>	Pyraclostrobin	<b>Empress® Intrinsic®</b> brand fungicide	12	G,N
		Fungal leaf spots, Rusts, Powdery mildew, <i>Rhizoctonia</i>	Trifloxystrobin	Compass®	12	G,N,L
	<b>Imidazolinone</b>	<i>Pythium</i> , <i>Phytophthora</i> , Downy mildew	Fenamidone	FenStop™	12	G
12	<b>Phenylpyrrole</b>	Fungal leaf spots, <i>Botrytis</i> , <i>Cylindrocladium</i> , <i>Fusarium</i> , <i>Rhizoctonia</i>	Fludioxonil	Medallion®, Emblem™	12	G,N,L
14	<b>Aromatic Hydrocarbon</b>	<i>Rhizoctonia</i>	PCNB	Revere®, Terraclor®	12	G,N, L
	<b>1, 2, 4-thiadiazole</b>	<i>Pythium</i> , <i>Phytophthora</i>	Etridiazole	Truban®, Terrazole®	12	G,N,L
17	<b>Hydroxyanilide</b>	<i>Botrytis</i>	Fenhexamid	Decree®	4	G,N
19	<b>Peptidyl Pyrimidine Nucleoside</b>	<i>Botrytis</i> , <i>Rhizoctonia</i> , <i>Alternaria</i> , Downy mildew, Powdery mildew, Anthracnose, Scab	Polyoxin-D	Affirm®	4	G,N,L
21	<b>Cyano-imidazole</b>	<i>Phytophthora</i> , <i>Pythium</i> , Downy mildew	Cyazofamid	Segway® Fungicide SC Segway® O	12	G,N,L
28	<b>Carbamate</b>	<i>Phytophthora</i> , <i>Pythium</i>	Propamocarb	Banol®	24	G,N,L
33 (Reclas- sified P07)	<b>Ethyl-phosphonate</b>	<i>Phytophthora</i> , <i>Pythium</i> , Downy mildew	Fosetyl-al (aluminium tris)	Aliette®, Areca	12	G,N,L
		<i>Phytophthora</i> , <i>Pythium</i> , Downy mildew	Phosphorous acid and salts	Agri FOS®, Alude®, Bio-phos®, Magellan®, Resyst®	4	G,N,L
40	<b>Cinnamic Acid Amides</b>	<i>Phytophthora</i> , Downy mildew	Dimethomorph	<b>Stature® SC</b> fungicide	12	G, N
	<b>Mandelic Acid Amides</b>	<i>Phytophthora</i> , Downy mildew	Mandipropamid	Micora®	4	G,N,L
41	<b>Tetracycline Antibiotic</b>	Bacterial diseases, Lethal yellows, Elm phloem necrosis	Oxytetracycline	Mycobject®, Tree Tech OTC®	NOL	L
43	<b>Pyridinylmethyl Benzamide</b>	<i>Phytophthora</i> , <i>Pythium</i> , Downy mildew	Fluopicolide	Adorn®	12	G,N,L
49	<b>Piperidinyl-thiazole-isoxazoline</b>	<i>Phytophthoras</i> , Downy mildew	Oxathiapiprolin	Segovis®	4	G,N,L

G = Greenhouse, N= Nursery, L=Landscape



## Fungicide Resistance Management Guide – (continued)

FRAC Group	Chemical Group(s)	Primary Target Disease(s)	Active Ingredient(s)	Fungicide Trade Name(s)	REI	Use Site(s)
M1	Copper, Complex	Fungal leaf spots, Bacterial diseases, Powdery mildew, Scab, Downy mildew	Copper sulfate	Camelot®	12	G,N,L
			Copper sulfate pentahydrate	Phyton® 27	24	G,N,L
	Copper, Fixed	Fungal leaf spots, Bacterial diseases, Powdery mildew, Scab, Downy mildew	Copper hydroxide	Kocide 2000®	24	G
M4	Phthalimide	Fungal leaf spots, <i>Botrytis</i> , Downy mildew, <i>Thielaviopsis</i>	Captan	Captan 50W	96	G,N,L
M5	Chloronitrile	Fungal leaf spots, <i>Botrytis</i> , Powdery mildew, Rusts, Scab, <i>Fusarium</i>	Chlorothalonil	Agrisel®, Chlorostar®, AllPro®, Concorde®, Echo®, Manicure®, Thaloni®, Daconil® Ultrex®, Legend®	12-48	G,N,L
M3	Dithio-carbamate	Fungal leaf spots, <i>Botrytis</i> , Rusts, Downy mildew, Scab	Mancozeb	Dithane®, Fore®, Formec®, Mancozeb®, Junction®, Pentathlon®, Protect®	24	G,N,L
		Fungal leaf spots, <i>Botrytis</i> , Rusts, Downy mildew	Ferbam	Ferbam Granuflo®	24	G, N
		Fungal leaf spots, <i>Botrytis</i> , Rusts	Ziram	Ziram Granuflo®	48	G, N
M3+3	Dithio-carbamate + Triazole	Fungal leaf spots, <i>Botrytis</i> , Rusts, Downy mildew, Powdery mildew	Mancozeb + Myclobutanil	Manhandle®	24	G,N,L
M5+3	Chloronitrile + Pyrimidine	Fungal leaf spots, <i>Botrytis</i> , Powdery mildew, Rusts, Scab	Chlorothalonil + Fenarimol	TwoSome™	NOEL*	L
	Chloronitrile + Triazole	Leaf spots and blights, Powdery mildew, Rusts, Scab, Flower blights	Chlorothalonil + Propiconazole	Concert® II	12	L
1+M3	Thiophanate + Dithio-carbamate	Fungal leaf spots, Powdery mildew, Downy mildew, <i>Botrytis</i>	Thiophanate-methyl + Dithiocarbamate	Zyban®	12	G,N,L
1+M5	Thiophanate + Chloronitrile	Fungal leaf spots, Powdery mildew, Downy mildew, <i>Botrytis</i>	Thiophanate-methyl + Chlorothalonil	ConSyst®, Spectro90®	12	G,N,L
1+2	Thiophanate + Dicarboxamide	Fungal leaf spots, <i>Botrytis</i> , <i>Rhizoctonia</i>	Thiophanate-methyl + iprodione	26/36®	24	G,N,L
1+14	Thiophanate + 1, 2, 4-thiadiazole	<i>Rhizoctonia</i> , <i>Thielaviopsis</i> , <i>Pythium</i> , <i>Fusarium</i>	Thiophanate-methyl + etridiazole	Banrot®	12	G,N,L
3+11	Triazole + Strobilurin	Rust, <i>Rhizoctonia</i> , <i>Botrytis</i> , Powdery mildew	Triadimefon + Trifloxystrobin	Trigo™	12	G,N,L

G = Greenhouse, N= Nursery, L=Landscape

1 Some products within the same chemical group may not control all pests listed, or there may be some products within the group that offer superior performance. Consult your local manufacturer's representative or your university/extension specialist if needed.

2 Specific use sites may vary by formulation and manufacturer; therefore, always read and consult product label directions.

## Fungicide Resistance Management Guide – (continued)

FRAC Group	Chemical Group(s)	Primary Target Disease(s)	Active Ingredient(s)	Fungicide Trade Name(s)	REI	Use Site(s)
4+12	<b>Acylalanine + Phenylpyrrole</b>	<i>Phytophthora</i> , <i>Pythium</i> , <i>Fusarium</i> , Crown and Stem rots	Mefenoxam + Fludioxonil	Hurricane®	48	G
7+11	<b>Pyrazole-4-Carboxamides + Strobilurin</b>	Fungal leaf spots, Downy mildew, Powdery mildew, Anthracnose, <i>Fusarium</i> , <i>Rhizoctonia</i> , <i>Phytophthora</i> , <i>Pythium</i> , Scab and Rusts, <i>Botrytis</i> , Blights	Benzovindiflupyr + Azoxystrobin	Mural®	12	G,N,L
		Fungal leaf spots, Downy mildew, Powdery mildew, Anthracnose, <i>Fusarium</i> , <i>Rhizoctonia</i> , <i>Phytophthora</i> , <i>Pythium</i> , Scab and Rusts, <i>Botrytis</i> , Blights, <i>Theleviopsis</i> spp.	Fluxapyroxad + Pyraclostrobin	<b>Orkestra® Intrinsic®</b> brand fungicide	12	G,N,L
	<b>Pyridine-Carboxamide + Strobilurin</b>	Fungal leaf spots, Downy mildew, Powdery mildew, Anthracnose, <i>Fusarium</i> , <i>Rhizoctonia</i> , <i>Phytophthora</i> , <i>Pythium</i> , Scab and Rusts, <i>Botrytis</i> , Blights	Boscalid + Pyraclostrobin	<b>Pageant® Intrinsic®</b> brand fungicide	12	G,N,L
	<b>Pyridinyl-ethyl-benzamides + Strobilurin</b>	Fungal leaf spots, Downy mildew, Powdery mildew, Anthracnose, <i>Fusarium</i> , <i>Rhizoctonia</i> , Scab and Rusts, <i>Botrytis</i> , Blights	Fluopyram + Trifloxystrobin	Broadform	12	G,N,L
9+12	<b>Anilinopyrimidine + Phenylpyrrole</b>	<i>Botrytis</i> , Powdery mildew, <i>Cercospora</i> , <i>Alternaria</i> , <i>Septoria</i> , <i>Myrothecium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>	Cyprodinil + Fludioxonil	Palladium	12	G,N
45+40	<b>Triazolo-Pyrimidylamine + Cinnamic Acid Amides</b>	Downy mildew, <i>Phytophthora</i>	Ametoctradin + Dimethomorph	Orvego® fungicide	12	G,N
FRAC Group	IRAC Group	Combination Fungicide / Insecticides Designated Within Fungicide Resistance Action Committee and Insecticide Resistance Action Committee Groups				
1	6	<b>Benzimidazole + Glycoside</b>	Vascular Pathogens: Wilts, Cankers, Dutch elm disease, <i>Fusarium</i>	Debacarb + Abamectin	Abasol®	NOL L
1	4A	<b>Benzimidazole + Neonicotinoid</b>	Vascular Pathogens: Wilts, Cankers, Dutch elm disease, <i>Fusarium</i>	Debacarb + Imidacloprid	Imisol®	NOL L

G = Greenhouse, N= Nursery, L=Landscape

## Fungicide Resistance Management Guide – (continued)

FRAC Group	Chemical Group(s)	Primary Target Disease(s)	Active Ingredient(s)	Fungicide Trade Name(s)	REI	Use Site(s)
44	<b>Biopesticide</b>	Botrytis, Powdery and Downy mildews, <i>Rhizoctonia</i> , <i>Pythium</i> , <i>Fusarium</i>	<i>Bacillus amyloliquefaciens</i> MBI 600	Serifel® NG fungicide	4	G,N
			<i>B. amyloliquefaciens</i> strain D747	Triathion® BA	4	G,N
		<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>	<i>Bacillus subtilis</i> GB03	Cease®	4	G,N
			<i>Bacillus subtilis</i> QST713	Rhapsody® AS	4	G,N,L
NC	<b>Botanical Extract</b>	Powdery mildew	Jojoba oil	Permatrol®	4	G,N,L
		Powdery mildew, Fungal leaf spots, Rusts, Scab	Clarified hydrophobic neem oil	Triact 70®	4	G,N,L
		Powdery mildew, Fungal leaf spots,	Macheaya extract	Qwel®	12	G
		Powdery mildew	<i>Raynouria sachalinesis</i>	Milsana®	24	G
PO5	<b>Botanical Extract</b>	Fungal and Bacterial leaf spots, Powdery mildew, Botrytis Gray Mold, Stem blights and rots	Extract of <i>Reynouria sachalinesis</i>	Regalia®	4	G,N,L
PO7	<b>Ethyl-phosphonate</b>	<i>Phytophthora</i> , <i>Pythium</i> , Downy mildew	Fosetyl-al	Aliette®	12	G,N,L
			Phosphorous acid	Agri FOS®, Alude®, ArborFox®, Bio-phos®, Magellan®, Resyst®	4	G,N,L
BM01	<b>Biopesticide</b>	<i>Phytophthora</i> , <i>Pythium</i> , <i>Fusarium</i> , <i>Cylindrocladium</i> , <i>Thielaviopsis</i>	<i>Trichoderma harzianum</i> T-22 + <i>T. virens</i> G-41	Rootshield® Plus (BW240)	4	G,N,L
			<i>Trichoderma harzianum</i> T22	PlantShield® HE, RootShield®	0	G,N,L
NC	<b>Biopesticide</b>	<i>Pythium</i> , <i>Rhizoctonia</i> , <i>Fusarium</i>	<i>Gliocadium virens</i> GL21	SoilGard® G	0	G,N,L
			<i>Streptomyces lydicus</i> WYEC108	Actinovate® , Actiono Iron	4	G,N,L
	<b>Hydrogen Dioxide/Peroxide</b>	Bacteria and Soilborne Fungi, Fungal leaf spots, Powdery mildew, Botrytis	Hydrogen Dioxide/Peroxide/Peroxyacetic Acid	ZeroTol®, TerraClean®, X3	0-2	G,N,L
	<b>Oil</b>	Powdery mildew	Unsulfonated residue of petroleum oil	SuffOil-X®	4	G,N
			Powdery mildew and Rusts	Mineral oil	Ultra-Pure Oil®	0
	<b>Quaternary Ammonium</b>	Bacteria/Algae	Quaternary amines	Green-Shield II®	0	G
			Quaternary ammonium chloride	KleenGrow®	48	G
<b>Soap</b>	Powdery mildew	Potassium salts of fatty acids	M-Pede®	12	G,N,L	

G = Greenhouse, N= Nursery, L=Landscape

1 Some products within the same chemical group may not control all pests listed, or there may be some products within the group that offer superior performance. Consult your local manufacturer's representative or your university/extension specialist if needed.

2 Specific use sites may vary by formulation and manufacturer; therefore, always read and consult product label directions.

**Fungicide Resistance Action Committee (FRAC) Mode of Action Classification List:**

- 1 Inhibition of tubulin formation in mitosis
- 2 Affect cell division, DNA and RNA synthesis and metabolism
- 3 DMI (Demethylation Inhibitor): Inhibition of sterol synthesis
- 4 Phenylamides - Affect RNA synthesis
- 5 Inhibition of an isomerase in sterol biosynthesis - Piperadines, Morpholines
- 7 Complex II in the mitochondria: succinate-dehydrogenase enzyme inhibitor
- 9 Affects amino acids and protein synthesis
- 11 Complex III: cytochrome bc1 (ubiquinol oxidase) at Qo site- Quinone outside- (cyt b gene) affecting respiration
- 12 MAP protein kinase in osmotic signal transduction
- 14 Lipid peroxidation
- 17 3-keto reductase during C4 demethylation in sterol biosynthesis
- 19 Chitin synthase
- 21 Complex III: cytochrome bc1(ubiquinone reductase) at Qi- Quinone inside- site in the mitochondria affecting respiration
- 28 Affects cell membrane permeability (proposed)
- 33 (Reclassified PO7)
- 40 Affects cell wall biosynthesis - cellulose synthase specifically
- 41 Protein synthesis attachment of aminoacyl-tRNA to ribosomal acceptor (A) site
- 43 Affects spectrin-like proteins, believed to play a role in maintaining the membrane stability
- 45 Complex III: cytochrome bc1 (ubiquinone reductase) at Qo site, stigmatellin binding sub-site affecting respiration
- U15 Oxysterol binding protein (OSBP) inhibition (proposed)
- M Multi-site contact activity
- NC Not classified

**Insecticide Resistance Action Committee (IRAC) Mode of Action Classification List:**

- 4a Neonicotinoids
- 6 Chloride channel activators

*The above lists are from the U.S. Environmental Protection Agency, in cooperation with the Fungicide Resistance Action Committee (FRAC) and the Insecticide Resistance Action Committee (IRAC) respectively.*

*FRAC and IRAC are technical working groups within the Global Crop Protection Federation (GCPF).*

*More information on the Fungicide Resistance Action Committee and the Mode of Action Classification is available from [www.frac.info](http://www.frac.info).*

*More information on the Insecticide Resistance Action Committee and the Mode of Action Classification is available from [www.irac.info](http://www.irac.info).*











# Mite and Insect Management

The BASF mite and insect management portfolio, which includes specialty insecticides, miticides and biological insect controls, provides solutions for both indoor and outdoor grower operations. Ask us about our innovative total release aerosol delivery system of insecticides that treats greenhouses of all sizes.

## Insecticide Use Sites

Product	Greenhouse	Container Nursery	Field Production	Conifer Nursery Plantation	Retail Greenhouse / Nursery	Interiorscape	Landscape and Recreational Areas	Vegetable Transplants and Production
<b>Specialty Insecticides</b>								
<b>Ventigra™</b> insecticide	■	■	■	■	■	■	■	■
<b>Sultan®</b> Miticide	■	■	■	■	■	■	■	
<b>Pylon®</b> Miticide/Insecticide	■							■
<b>DuraGuard® ME</b> Microencapsulated Insecticide	■	■	■				■	
<b>Siesta®</b> Fire Ant Bait		■	■	■	■		■	
<b>Amdro® Pro</b> Insect Fire Ant Bait	■	■	■	■	■		■	
<b>Ultra Pure Oil®</b> Horticultural Insecticide, Miticide and Fungicide	■	■	■	■	■	■	■	■
<b>Biological Insect Controls</b>								
<b>Nemasys®</b> Beneficial Nematodes	■	■	■	■	■	■		
<b>Millenium®</b> Beneficial Nematodes	■	■	■	■	■	■	■	
<b>Velifer®</b> Fungal Contact Insecticide	■							■
<b>Total Release Insecticides</b>								
<b>Beethoven® TR</b> Miticide/Insecticide	■							
<b>Pylon TR</b> Total Release Miticide/Insecticide	■							
<b>1300 Orthene® TR</b> Total Release Insecticide	■							
<b>Attain® TR</b> Total Release Insecticide	■							
<b>Duraplex® TR</b> Total Release Insecticide	■							
<b>Pyrethrum® TR</b> Total Release Insecticide	■							



# Ventigra™

## Insecticide

**Ventigra** insecticide provides effective control of piercing-sucking insect pests in production and landscape ornamentals and vegetable transplants for the home consumer market. Ventigra Insecticide is a unique mode of action subgroup with no known cross-resistance with key commercial insecticide classes. Ventigra Insecticide has minimal impact to beneficial insects including predatory mites, parasitoid wasps, and pollinators, and it's compatible with biological releases in a production program.

**IRAC group:** 9D

**Active ingredient:**

Afidopyropen

**Signal word:** CAUTION

**Formulation:**

DC (Dispersible Concentrate)

**Use rates:** 1.4 - 7.0 fl oz per 100 gal I

**PPE:** Long sleeved shirt and long pants, chemical-resistant gloves made of barrier laminate or butyl rubber or neoprene rubber, shoes and socks

**REI:** 12 Hours

**Packaging:** 4 x 20 fl oz

### Key Features and Benefits

- Rapid onset of action to stop insect feeding and damage
- Long residual control
- Translaminar movement delivers optimal protection against actively feeding pests
- No negative impact on bee populations, including colony health and development
- Excellent rotational partner in IPM programs
- Unique MOA subgroup controls pests that have developed resistance to existing insecticides

### Insects Controlled

- Aphids, whiteflies, and mealybug, and has suppression for certain species of scale.

### Use Sites

- Commercial and retail nurseries, field and container
- Commercial, public, and recreational gardens
- Commercial and residential landscapes, including golf courses, parks and recreation areas
- Greenhouses, shadehouses, and lathhouses
- Vegetable transplants
- Interiorscapes
- Forest tree and conifer nurseries and plantations

Labeled for use on juvenile fruit and nut trees, vines, brambles and bushberries grown in commercial ornamental production nurseries.

Labeled for vegetable transplants for the home consumer market and vegetables grown in commercial greenhouses.



# Sultan®

## Miticide

**Sultan** miticide is for the grower who wants fast, powerful control and lasting protection against all life stages of spider mites in greenhouse and nursery crops.

### Key Features and Benefits

- Fast knockdown—typically within three hours of contact
- Controls mite populations for up to 28 days
- Unique mode of action increases rotation options
- Compatible with Integrated Pest Management (IPM) programs using beneficial insects and predatory mites
- May be used in interiorscapes and outdoor landscape settings

### Mites Controlled

- Two-spotted spider mite
- Glover mite
- Pacific spider mite
- Carmine mite
- Citrus red mite
- Spruce spider mite

### Use Sites

- Greenhouses
- Lathhouses and shadehouses
- Interiorscapes
- Commercial and retail nurseries (field or container)
- Commercial, public and recreational gardens
- Landscape ornamentals
- Commercial and residential landscape maintenance areas
- Tree and conifer nurseries or plantations



**IRAC group:** 25

**Active ingredient:**  
Cyflumetofen

**Signal word:** CAUTION

**Formulation:**  
Suspension

**Use rates:** 13.7 fl oz  
per 100 gal

**PPE:** Chemical-resistant gloves, long-sleeved shirt, long pants, shoes plus socks

**REI:** 12 Hours

**Packaging:** 4 x 16 fl oz  
per case





**IRAC group:** 13

**Active ingredient:**

Chlorfenapyr

**Signal word:** CAUTION

**Formulation:**

Suspension

**Use rates:** 2.6 -10.0 fl oz per 100 gal

**PPE:** Coveralls, chemical-resistant gloves, shoes plus socks

**REI:** 12 hours

**Packaging:** 4 x 16 fl oz per case; 2 x 0.5 gal per case

# Pylon®

Miticide/Insecticide

**Pylon** miticide/insecticide is the greenhouse grower's most effective treatment for control of Western flower thrips, chilli thrips and mites.

**Pylon** miticide/insecticide is also available as **Pylon TR**, a total release formula, in the Total Release section of this guide.

## Key Features and Benefits

- Broad-spectrum control of nymph and adult stages of mites
- The foundation for an effective thrips control program
- Translaminar activity maximizes distribution within the treated plants
- Excellent finish treatment to achieve the crop quality expectations of the most discerning customers

## Key Insects Controlled

- Western Flower Thrips
- Chilli Thrips
- Spider Mites
- Broad Mite
- Rust Mite
- Cyclamen Mite
- Citrus Budmite
- Fungus Gnats
- Caterpillars
- Foliar Nematodes

## Use Sites

- Commercial ornamental greenhouses
- Commercial production greenhouses for fruiting vegetables



# DuraGuard® ME

Microencapsulated Insecticide

**DuraGuard ME** is for the production ornamental grower who wants long-lasting control of a wide range of pests.

## Key Features and Benefits

- Highly versatile, long-lasting residual
- Micro-encapsulated technology
- Increased plant safety, even on many open blooms
- Little to no odor

## Key Insects Controlled

- |                      |                |
|----------------------|----------------|
| ■ Aphids             | ■ Leafhoppers  |
| ■ Fungus gnats       | ■ Leaf miners  |
| ■ Fungus gnat larvae | ■ Flea beetles |
| ■ Mealybugs          | ■ Thrips       |
| ■ Scale insects      | ■ Ants         |
| ■ Shore flies        | ■ Crickets     |
| ■ Spider mites       |                |

## Use Sites

- Commercial ornamental greenhouses and nurseries
- Landscaped outdoor areas



### IRAC group: 18

Restricted Use Pesticide

### Active ingredient:

Chlorpyrifos

**Signal word:** CAUTION

### Formulation:

ME (microencapsulated liquid concentrate)

### Use rates:

*Foliar Spray:* 25 - 50 fl oz per 100 gal

*Drench:* 0.125 fl oz per gal

*Outdoor Ornamentals:* 0.5-1.75 fl oz per 1000 sq ft

**PPE:** Long-sleeved shirt and long pants, waterproof gloves, shoes plus socks

**REI:** 24 hours

**Packaging:** 4 x 1 gal





**IRAC group:** 22B

**Active ingredient:**  
Metaflumizone

**Signal word:** CAUTION

**Formulation:**  
Granular bait

**Use rates:**  
*Broadcast treatment:*  
1.0-1.5 lb per acre

*Mound treatment:*  
2-4 level tablespoons

**PPE:** Long-sleeved shirt  
and long pants,  
waterproof gloves,  
shoes plus socks

**REI:**  
12 hours

**Packaging:** 15 lb bag

# Siesta®

## Insecticide Fire Ant Bait

**Siesta** insecticide fire ant bait is an improved baiting formulation that provides the professional horticulturalist with fast, long-lasting control of native and red imported fire ants.

### Key Features and Benefits

- Non-repellent, with a novel mode of action
- Eliminates the queen and the entire colony
- Rapid cessation of feeding leads to colony collapse in three days
- Listed on USDA APHIS for nursery stock RIFA quarantine certification

### Insects Controlled

- Native and red imported fire ants

### Use Sites

#### Ornamental sites:

- Outdoor nurseries (field and container)
- Soil around bearing and non-bearing fruit or nut trees

#### Non-cropland areas:

- Airports
- Cemeteries
- Parks
- School and picnic grounds
- Recreational areas
- Roadside areas

#### Turf sites:

- Athletic fields
- Commercial grounds
- Golf courses
- Lawns
- Landscaped areas
- Sod farms





# Amdro® Pro

## Insecticide Fire Ant Bait

**Amdro Pro** fire ant bait is a cost-effective mound baiting system for control of imported and native fire ants, and other nuisance ants

### Key Features and Benefits

- Eliminates the queen and colony within seven days of mound treatment
- Rids entire area of fire ants within two weeks of broadcast application
- Approved for aerial applications
- Listed by USDA APHIS for RIFA quarantine certification of nursery stock

### Key Insects Controlled

- Imported and native fire ants
- Little fire ants
- Harvester ants
- Bigheaded ants

### Use Sites

Ornamentals sites:

- Outdoor container nurseries
- Outdoor field nurseries
- Soil around non-bearing fruit or nut trees

Non-cropland areas:

- Airports
- Cemeteries
- Parks
- School and picnic grounds
- Recreational areas
- Roadside areas
- Pasture and rangeland

Turf sites:

- Athletic fields
- Commercial grounds
- Golf courses
- Lawns
- Landscaped areas
- Sod farms



**IRAC group:** 20A

**Active ingredient:**

Hydramethylnon

**Signal word:** CAUTION

**Formulation:**

Granular bait

**Use rates:**

*Broadcast treatment:*

1-2 lb per acre depending on ant species

*Mound treatment:*

2 to 5 level tablespoons

**PPE:** Long-sleeved shirt and long pants, chemical-resistant gloves, such as rubber or waterproof gloves, shoes plus socks

**REI:** 12 hours

(agricultural uses only)

**Packaging:** 25 lb bag





# Ultra Pure Oil

Horticultural Insecticide, Miticide and Fungicide

A flexible, year-round, 99% pure spray oil, **Ultra-Pure Oil** meets the pest and disease resistance management needs of horticulturalists by providing highly effective contact activity with lower potential for plant phytotoxicity.

## Key Features and Benefits

- Broad-spectrum control of insects, mites and fungi
- Use alone or in tank-mixes
- No odor and no post-harvest restrictions
- Excellent disease resistance management tool

## Pests Controlled

- Adelgids
- Aphids
- Fungus gnats
- Lacebugs
- Leaf miners (larvae)
- Leafhoppers
- Mealybugs
- Mites
- Scale insects
- Thrips
- Whiteflies
- Webworms
- Powdery mildew on ornamentals

## Use Sites

- Greenhouses
- Nurseries
- Container gardens
- Bedding plants
- Christmas tree farms
- Ornamental trees and shrubs
- Flowering and foliage plants
- Field crops
- Fruit and nut crops
- Vegetables and herbs

**IRAC group:** Horticultural oil

**Active ingredient:**

- 98% petroleum oil

**Signal word:** CAUTION

**Formulation:**

Liquid

**Use rates:** 0.5 gal to 2 gal per 100 gal

**PPE:** Long-sleeved shirt and long pants, chemical-resistant gloves, shoes plus socks

**REI:** 4 hours

**Packaging:**

2 x 2.5 gal, 30 gal drum



# Nemasys®

## Beneficial Nematodes

**Nemasys** beneficial nematodes help the professional greenhouse grower achieve biocontrol of fungus gnats and Western flower thrips. No re-entry restrictions means fewer worker interruptions and optimizes production efficiencies.

### Key Features and Benefits

- Compatible with other biologicals and many common chemistries including fertilizers, fungicides and insecticides
- Excellent resistance management tool
- Foundation of biological control programs

### Key Insects Controlled

- Fungus Gnats
- Western flower thrips

### Use Sites

- Commercial greenhouses
- Interiorscapes
- Field, container and retail nurseries



**Active ingredient:**  
*Steinernema feltiae*

**PPE:** None required.

**REI:** Zero. No re-entry restrictions.

**Packaging:** 50 million nematode per tray, or 5 x 250 million nematodes per case





**Active ingredient:**

*Steinernema carpocapsae*

**PPE:** None required.

**REI:** Zero. No re-entry restrictions.

**Packaging:** 250 million nematodes per tray

# Millenium®

## Beneficial Nematodes

**Millennium** beneficial nematodes help the ornamental grower achieve effective biocontrol of shore fly pests in greenhouses, with zero re-entry level use restrictions.

### Key Features and Benefits

- Compatible with other biologicals and many common chemistries including fertilizers, fungicide and insecticides
- Excellent resistance management tool
- Foundation of biological control programs

### Key Insects Controlled

- Shore flies

### Use Sites

- Commercial greenhouses
- Interiorscapes
- Field, container and retail nurseries
- Lawns and landscapes





# Velifer®

## Fungal Contact Insecticide/Miticide

**Velifer** fungal contact insecticide/miticide effectively controls major greenhouse pests, including whiteflies, and is an invaluable tool for growers to use alone or in rotation with chemicals to reduce resistance and extend the life of their chemicals.

### Key Features and Benefits

- Liquid oil dispersion formulation
- Biological product based on a naturally occurring beneficial fungus
- Active against all life stages of the target pest
- Works in a manner that complements chemical solutions
- Resistance development is highly unlikely
- No known adverse symptoms of phytotoxicity
- Does not affect beneficial insects

### Key Insects Controlled

- Aphid
- Whitefly
- Mealybug
- Thrips
- Mites

### Use Sites

- Commercial greenhouses



### Active ingredient:

*Beauveria bassiana* strain  
PPRI 5339

**Signal word:** CAUTION

### Formulation:

Liquid oil dispersion

### Use rates:

3 - 13 fl oz per 100 gal water

**PPE:** Long-sleeved shirt,  
long pants, shoes and  
socks, protective eyewear

**REI:** 4 hours

**Packaging:**



# Insecticide Resistance Management Guide

IRAC Group	Chemical Sub Group(s)	Primary Target Insect(s)	Active Ingredient(s)	Insecticide Trade Name(s)	REI	Use Site(s)
1A	<b>Carbamates</b>	Caterpillars, Soft scale insects, Thrips*	Carbaryl	Sevin®	12	N,L
			Methiocarb	Mesuro®	24	G,N
1B	<b>Organophosphates</b>	Aphids, Caterpillars, Mealybugs*, Ants, Whiteflies (general and B-type), Fungus gnats, Shore flies, Glassy-winged sharpshooter	Acephate	<b>1300 Orthene® TR</b> total release insecticide, <b>Orthene® TT&amp;O</b> WSP insecticide, Pinpoint® PT 1300, <b>Tame®/ Orthene TR</b> total release insecticide	24	G,N,L
			Chlorpyrifos	<b>DuraGuard® ME</b> microencapsulated insecticide, <b>DuraPlex® TR</b> total release insecticide, Dursban	24	G,N
			Diazinon	Diazinon®	12	N
			Naled	Diabrom®	24	G
			Dimethoate	Dimethoate® 267, Cygon® 2E, Dimate® 4EC	48	N
			Malathion	Malathion® 8F	12	N,L
			Methidathion	Supracide®	48	N
			Oxydemeton-methyl	MSR® Spray Concentrate	48-72	N
			Phosmet	Imidan® 70 W	24	N,L
			Trichlorfon	Dylox® 80	N/A	L
2A	<b>Cyclodiene Organochlorides</b>	Aphids, Mites, Borers, True bugs and Whiteflies	Endosulfan	Fulex Thiodan® insecticidal smoke, Thionex® EC	24	G
2	<b>Phenylpyrazoles (Fiproles)</b>	Imported fire ants, Nuisance ants, Fleas, Ticks, Mole crickets	Fipronil	Ceasefire®, Topchoice®	24	L
3	<b>Pyrethroids</b>	Aphids, Caterpillars, Glassy-winged sharpshooter, Mealybugs*, Thrips*, Scales*	Bifenthrin	OnyxPro®, Talstar®, <b>Attain TR®</b> total release insecticide	12	G,N,L
			Cyfluthrin	Decathlon®, Tempo®	12	G,N,L
			Fenpropathrin	Tame®	24	G,N,L
			Fluvalinate	Mavrik® AF	12	G,N,L
			Lambda-cyhalothrin	Scimitar® GC	24	G,N,L
			Permethrin	Astro®, Perm-Up®, Permethrin® 3.2 EC	12	L
			Deltamethrin	DeltaGard®	12	N,L
	<b>Pyrethrins</b>	Aphids, Caterpillars, Glassy-winged sharpshooter, Mealybugs*, Thrips*, Scales*	Pyrethrins	<b>Pyrethrum® TR</b> total release insecticide	12	G,N

G = Greenhouse, N= Nursery, L=Landscape

\* Some, but not all, species of selected pests within the group are controlled. Read the product specimen label for additional information. Consult your local university/extension specialist if needed.

1 Some products within the same chemical group may not control all pests listed, or there may be some products within the group that offer superior performance.

2 Specific use sites may vary by formulation and manufacturer; therefore, always read and consult product label directions.

## Insecticide Resistance Management Guide – (continued)

IRAC Group	Chemical Sub Group(s)	Primary Target Insect(s)	Active Ingredient(s)	Insecticide Trade Name(s)	REI	Use Site(s)
4A	<b>Neonicotinoids</b>	Aphids, Whiteflies (general, B- and Q-type), Grubs*, Scales*, Mealybugs & Scale, Caterpillars, Midges, Thrips	Acetamiprid	TriStar®	12	G,N,L
			Dinotefuran	Safari®	12	G,N,L
			Imidacloprid	Marathon®, Merit®, Discus® Tablets	0-12	G,N,L
			Imidacloprid + Cyfluthrin	Discus® NG	12	N
			Imidacloprid + Bifenthrin	Allectus® G	12	L
			Imidacloprid + Deba carb	Imisol®	NOL	L
			Imidacloprid + 12-9-4 fertilizer carrier	Coretect®	NOL	L
			Imidacloprid slow release	SusCon®	NOL	L
			Bifenthrin + Clothianidin	Allectus®, Aloft®	12	L
		Dinotefuran	Safari®	12	G,N,L	
		Thiamethoxam	Flagship 25 WG®	12	G,N	
4B	<b>Nicotine</b>	Whiteflies, Spider mites, Aphids, Mealybugs	Nicotine	Fulex® Nicotine® Fumigator,* Nicotine Smoke Generator		G
5	<b>Spinosyns</b>	Caterpillars, Mites*, Thrips	Spinosad	Conserve®	4	G,N,L
		Caterpillars, Leafminers, Red Imported Fire Ants	Spinosad	Entrust® SC	4	G,N
6	<b>Avermectins, Milbemycins</b>	Mites, Thrips	Abamectin	Avid®	12	G,N,L
			Abamectin + Deba carb	Abasol®	NOL	L
	Mites	Milemectin	Ultiflora Miticide®	12	N	
	<b>Glycoside</b>	Caterpillars, Leafminers, Mites, Psylla	Emamectin Benzoate	Enfold®	12	N
7A		Whiteflies, Fungus gnats, Scales, Mealybugs, Aphids	s-Kinoporone	Enstar® AQ	4	G
	<b>Juvenile Hormone Analogues</b>	Imported fire ants	Methoprene	Extinguish®	4	N,L
7B	<b>Fenoxycarb</b>	Ants	Fenoxycarb	Award®	12	G
7C	<b>Pyriproxyfen</b>	Fungus gnats, Shore flies	Pyriproxyfen	Distance®, Fulcrum®	12	G,N
9B	<b>Pyridine Azomethine Derivatives</b>	Aphids, Mealybugs*, Whiteflies*	Pymetrozine	Endeavor®	12	G,N,L
10A	<b>Clofentezine, Hexythiazox</b>	Mites	Clofentezine	Ovation®, Applause™,	12	G,N
			Hexythiazox	Hexygon DF®	12	G,N,L

G = Greenhouse, N= Nursery, L=Landscape

## Insecticide Resistance Management Guide – (continued)

IRAC Group	Chemical Sub Group(s)	Primary Target Insect(s)	Active Ingredient(s)	Insecticide Trade Name(s)	REI	Use Site(s)
10B	<b>Etoxazole</b>	Mites*	Etoxazole	TetraSan® <b>Beethoven TR</b> ® miticide/insecticide	4-24	G,N,L G
11A	<b>Bacillus thuringiensis and the insecticidal proteins they produce</b>	Fungus gnat larvae, Caterpillars	<i>Bacillus thuringiensis</i> sub <i>Israelensis</i>	Gnatrol®, Deliver®, Javelin® WG	4	G,N,L
12B	<b>Organotin Miticides</b>	Mites*	Fenbutatin-oxide	ProMITE® 50 WP	48	G,N,L
13	<b>Pyrroles</b>	Mites*, Thrips, Fungus gnats, Foliar nematodes, Caterpillars	Chlorfenapyr	<b>Pylon</b> ® Insecticide, <b>Pylon</b> ® TR Insecticide	12 4- 12	G G,N
15	<b>Benzoylureas</b>	Fungus gnats	Diflubenzuron	Adept®	12	G
		Fungus gnats	Diflubenzuron	Dimilin® SC	12	G,N
		Aphids, Mites, Thrips	Novaluron	Pedestal®	12	G,N
16	<b>Buprofezin</b>	Mealybugs*, Whiteflies*	Buprofezin	Talus®	12	G,N,L
17	<b>Cyromazine</b>	Fungus gnats, Leafminers	Cyromazine	Citation®	12	G,N,L
18	<b>Diacylhydrazines</b>	Caterpillars	Tebufenozide	Confirm®	4	N
			Methoxyfenozide	Intrepid®	4	G,N
20A	<b>Hydramethylnon</b>	Imported fire ants	Hydramethylnon	<b>Amdro Pro</b> ® fire ant bait insecticide	12	N,L
20B	<b>Acequinocyl</b>	Mites	Acequinocyl	Shuttle® O	12	G,N,L
21A	<b>Pyridazinones</b>	Mites, Whiteflies* (general and Q-type)	Pyridaben	Sanmite®	12	G,N
			Fenazaquin	Magus®	12	G,N
	<b>METI Acaricides and Insecticides</b>	Spider mites	Fenpyroximate	Akari®	12	G
22A	<b>Oxadizines</b>	Caterpillars, European crane fly, Ants including fire ants	Indoxacarb	Advion®	N/A	L
22B	<b>Semicarbazones</b>	Fire ants	Metaflumizone	<b>Siesta</b> ® fire ant bait insecticide	12	N,L
23	<b>Tetronic and Tetramic Acid Derivatives</b>	Spider Mites*, Whiteflies*	Spiromesifen	Judo®, Forbid®, Savate™	12	G,N
		Aphids, Leafhoppers, Mealybugs, Psyllids, Whiteflies, Mites	Spirotetramat	Kontos®	24	G,N

G = Greenhouse, N= Nursery, L=Landscape

\* Some, but not all, species of selected pests within the group are controlled. Read the product specimen label for additional information. Consult your local university/extension specialist if needed.

1 Some products within the same chemical group may not control all pests listed, or there may be some products within the group that offer superior performance.

2 Specific use sites may vary by formulation and manufacturer; therefore, always read and consult product label directions.

## Insecticide Resistance Management Guide – (continued)

IRAC Group	Chemical Sub Group(s)	Primary Target Insect(s)	Active Ingredient(s)	Insecticide Trade Name(s)	REI	Use Site(s)
25	<b>Beta-ketonitrile Derivatives</b>	Spider mites	Cyflumetofen	<b>Sultan</b> ® miticide	12	G,N,L
28	<b>Diamides</b>	Aphids, Whiteflies, Thrips	Cyantraniliprole	Mainspring® Mainspring® GNL	4	G,N,L
29	<b>Flonicamid</b>	Aphids, Glassy-winged sharpshooter, Mealybugs, Thrips	Flonicamid	Aria®	12	G,N,L
NC	<b>Oils</b>	Mites, Mealybugs, Aphids, Thrips	Clarified hydrophobic extract of neem oil	Triact 70®	4	G,N,L
			Petroleum oil	<b>Ultra-Pure Oil</b> ®, PureSpray®, Green Organic Horticultural Spray Oil	4	G,N,L
	<b>Soaps</b>	Mites*, Mealybugs*, Whiteflies*	Potassium salts of fatty acids	M-Pede®	12	G,N,L
	<b>Inorganics</b>	Slugs, Snails	Iron phosphate	Sluggo®	0	L
	<b>Aldehydes</b>	Slugs, Snails	Metaldehyde	Durham®, Trail's End®, Slug-It®	12	L
	<b>Biological Agents</b>	Nematodes*	<i>Myrothecium verrucaria</i>	DiTera®	4	N,L
		Fungus gnat	<i>Steinernema feltiae</i>	<b>Nemasys</b> ® beneficial nematodes	0	G,I,N
		Grubs	<i>Heterorhabditis bacteriophora</i>	<b>Nemasys G</b> ® beneficial nematodes	0	G,N
		Black vine weevil	<i>Steinernema kraussei</i>	<b>Nemasys L</b> ® beneficial nematodes	0	G,N
		Weevils*, Borers*, Shore flies	<i>Steinernema carpocapsae</i>	<b>Millenium</b> ® beneficial nematodes	0	G,N,L
Weevils*		<i>Steinernema carpocapsae</i>	<b>Nematac C</b> beneficial nematodes	0	G,N,L	
Aphids, Q-type Whitefly, Thrips		<i>Beauveria bassiana</i>	Naturalis® L, BotaniGard®	4	G,N,L	
UN	<b>Pyridalyl</b>	Thrips, Caterpillars	Pyridalyl	Overture®	12	G
	<b>Dicofol</b>	Mites*	Dicofol	Dicofol 4-E	12-48	G,N,L
	<b>Pyridine Azomethines</b>	Whiteflies, Aphids, Mealybugs	Pyrafluquinazon	Rycar®	12	G
	<b>Carbazates</b>	Spider mites	Bifenazate	Floramite®	4	G,N,L
	<b>Biopesticide Insect Growth Regulator</b>	Aphids, Thrips, Whiteflies	Azadirachtin	Azatin O®	4	G,N
	<b>Diacylhydrazines</b>	Aphids, Thrips*, Whiteflies*	Azadirachtin	Azatin XL®, Ornazin® Captiva®	4-12 4	G,N,L G,N
6+UN		Mites, Leaf Miners	Bifenazate + Abamectin	Sirocco™	12	G,N,L

G = Greenhouse, N= Nursery, L=Landscape



**Insecticide Resistance Action Committee (IRAC) Mode of Action Classification List:**

- 1 Acetylcholine esterase inhibitor
- 2 GABA-gated chloride channel antagonists
- 3 Sodium channel modulators
- 4 Nicotinic Acetylcholine receptor agonists/antagonists
- 5 Nicotinic Acetylcholine receptor agonists (allosteric) (not group 4)
- 6 Chloride channel activators
- 7 Juvenile hormone mimics
- 9 Compounds of unknown or non-specific mode of action (selective feeding blockers)
- 10 Compounds of unknown or non-specific mode of action (mite growth inhibitors)
- 11 Microbial disruptors of insect midgut membranes (includes transgenic crops expressing B.t. toxins)
- 12 Inhibitors of oxidative phosphorylation, disruptors of ATP formation (inhibitors of ATP synthase)
- 13 Uncouplers of oxidative phosphorylation via disruption of proton gradient
- 15 Inhibitors of chitin biosynthesis, type 0, Lepidopteran
- 16 Inhibitors of chitin biosynthesis, type 1, Homopteran
- 17 Ecdysone receptor agonists
- 18 Ecdysone agonists
- 20 Mitochondrial complex III electron transport inhibitors (Coupling site II)
- 21 Mitochondrial complex I electron transport inhibitors
- 22 Voltage-dependent sodium channel blocker
- 23 Inhibitors of lipid synthesis
- 25 Neuronal inhibitors (unknown mode of action)
- 28 Ryanodine receptor modulator
- 29 Chordotonal organ modulators – undefined target site
- NC Not classified by IRAC

*The above list is from the U.S. Environmental Protection Agency, in cooperation with the Insecticide Resistance Action Committee (IRAC). IRAC is a technical working group within the Global Crop Protection Federation (GCPF).*

*More information on the Insecticide Resistance Action Committee and the Mode of Action Classification is available from [www.irc.info](http://www.irc.info).*







# Weed Management

Control weeds with BASF pre- and post-emergent herbicides in granular and liquid formulations. Designed with plant safety in mind, BASF pre-emergent herbicides provide broad-spectrum weed control with outstanding results. Flexible use rates and application methods result in reduced hand weeding costs, resulting in more efficient production of the highest quality ornamental nursery crops.

## Herbicide Use Sites

Product	Bareground	Conifer Nursery Plantation	Container Nursery	Field Production	Retail Nursery	Landscape and Recreational Areas	Greenhouse and Shadehouse
<b>Preemergent Herbicides</b>							
<b>FreeHand® 1.75G</b> Herbicide	■	■	■	■	■	■	
<b>Pendulum® AquaCap</b> Herbicide	■	■	■	■	■	■	
<b>Pendulum® 3.3 EC</b> Herbicide	■	■	■	■	■	■	
<b>Pendulum® 2G</b> Herbicide	■	■	■	■	■	■	
<b>Tower®</b> Herbicide	■ <sup>1</sup>	■	■	■	■	■	■
<b>Postemergent Herbicides</b>							
<b>Finale®</b> Herbicide	■	■	■	■	■	■	■
<b>Basagran® T&amp;O</b> Herbicide	■	■	■	■	■	■	
<b>Segment® II</b> Herbicide	■	■	■	■	■	■	

<sup>1</sup> Tower herbicide label allows for use on gravel or ground floors of open-sided lathhouses (shadehouses) or other polyhouse structures that allow poly

## Weeds Controlled

Common name	Scientific name	Pendulum® herbicide products	FreeHand® 1.75G herbicide	Tower® herbicide	Finale® herbicide	Basagran® T&O herbicide	Segment® II herbicide
American burnweed	<i>Erechtites hieraciifolia</i>		■				
Barnyardgrass	<i>Echinochloa crus-galli</i>	■	■	■	■		■
Bittercress	<i>Cardamine</i> spp.	■	■	■			
Annual bluegrass	<i>Poa annua</i>	■	■	■	■		
Carpetweed	<i>Mollugo verticillata</i>	■	■	■			
Chamberbitter	<i>Pyllanthus urinaria</i>		■				
Common chickweed	<i>Stellaria media</i>	■	■		■		
Crabgrass	<i>Digitaria</i> spp.	■	■	■	■		■
Dogfennel	<i>Eupatorium capillifolium</i>		■				
Doveweed, Marsh dayflower	<i>Murdannia</i> spp.		■	■			
Eclipta	<i>Eclipta prostrata</i>		■	■			
Foxtail	<i>Setaria</i> spp.	■	■	■	■		■
Galinsoga	<i>Galinsoga</i> spp.		■	■		■	
Goosegrass	<i>Eleusine indica</i>	■	■	■	■		■
Common groundsel	<i>Senecio vulgaris</i>		■	■		■	
Henbit	<i>Lamium amplexicaule</i>	■	■				
Johnsongrass	<i>Sorghum halepense</i>	■		■	■		■
Common lambsquarters	<i>Chenopodium album</i>	■	■	■	■	■	
Liverwort	<i>Marchantia</i> sp.		■	■			
Longstalked phyllanthus	<i>Phyllanthus tenellus</i>		■				
Marsh parsley	<i>Apium leptophyllum</i>		■				
Mouse-ear chickweed	<i>Cerastium vulgatum</i>	■	■		■		
Mulberry weed	<i>Fatoua villosa</i>		■				
Panicum, various	<i>Panicum</i> spp.	■	■	■	■		■
Birdseye pearlwort	<i>Sagina procumbens</i>		■	■			
Pigweed	<i>Amaranthus</i> sp.	■	■	■	■		
Evening primrose	<i>Oenothera biennis</i>	■	■				
Common Purslane	<i>Portulaca oleracea</i>	■	■	■	■	■	
Florida pusley	<i>Richardia scabra</i>	■	■	■			
Sedges	<i>Cyperus</i> spp.		■	■	■	■	
Shepherd's purse	<i>Capsella bursa-pastoris</i>	■	■	■	■	■	
Pennsylvania smartweed	<i>Polygonum pennsylvanicum</i>	■	■		■	■	
Sowthistle	<i>Sonchus</i> spp.		■				
Spurge, various	<i>Euphorbia</i> spp. (formerly <i>Chamaesyce</i> spp.)		■	■			
Willowherb	<i>Epilobium</i> spp.			■			
Yellow woodsorrel	<i>Oxalis</i> spp.	■	■		■	■	
Tassel-flower	<i>Emilia</i> spp.	■	■				
Thistle	<i>Cirsium</i> spp.					■	
Field sandbur	<i>Cencherus spinifex</i>	■	■	■	■		■
Wild sunflower	<i>Helianthus</i> spp.					■	
Wild mustard	<i>Sinapis arvensis</i>				■	■	

This is not a complete list of weeds controlled. Consult the product label for weeds controlled.



# FreeHand® 1.75G

## Herbicide

For the nursery grower who wants the convenience of granular applications, **FreeHand 1.75G** herbicide delivers unsurpassed preemergent control of the toughest summer weeds.

### Key Features and Benefits

- Combines the power and efficacy of proven ingredients
- Effective control of annual grasses, sedges and small-seeded broadleaf weeds
- Unsurpassed performance on difficult to control weeds such as spurge, groundsel, yellow nutsedge, kyllinga and doveweed
- Controls more than 50 of the most troublesome weeds in nursery production

### Weeds Controlled

Refer to the *Weeds Controlled* chart for common weeds controlled. For a complete listing, visit [www.CDMS.net](http://www.CDMS.net) or find the product label using the convenient QR Code link below.

### Use Sites

- Commercial nurseries (container, liner, and field)
- Residential and commercial landscapes
- General grounds maintenance (noncrop areas)
- Conifer, Christmas tree and hardwood liner nurseries and tree plantations
- Turfgrass

May be applied to juvenile fruit and nut trees, vines, brambles and bushberries grown in commercial ornamental production nurseries. Always read and consult product label directions.



### WSSA group:

3 + 15

### Active ingredients:

Dimethenamid-P  
Pendimethalin

### Signal word: CAUTION

### Formulation: G (Granular)

### Use rates:

*Commercial ornamental production:* 100, 150, and 200 lb per acre

**PPE:** Coverall, waterproof gloves, shoes plus socks, protective eyewear

**REI:** 12 hours

**Packaging:** 50 lb bags





# Pendulum® AquaCap

Herbicide

**Pendulum AquaCap**, a microencapsulated formulation of pendimethalin, provides nursery growers with highly effective preemergent control of the toughest grassy and broadleaf weeds, and the confidence of proven plant safety.

## Key Features and Benefits

- Controls more than 40 of the most troublesome weeds in nursery production
- Virtually no odor
- Reduced potential for staining
- Ease and flexibility in handling, mixing and cleanup

## Weeds Controlled

Refer to the *Weeds Controlled* chart in this guide for common weeds controlled. For a complete listing, visit [www.CDMS.net](http://www.CDMS.net) or find the product label using the convenient QR Code link below.

## Use Sites

- Field-, liner-, and container-grown ornamental production
- Bare ground areas for container placement
- Bulb plantings
- Non-bearing fruit and nut tree nursery production
- Conifer and hardwood seedling nursery production
- Tree plantations for site preparation and maintenance
- Residential and commercial landscape
- Golf courses
- Recreational parks and athletic fields
- Sod farms
- General grounds maintenance (non-cropland areas)

**WSSA group:** 3

**Active ingredient:**

Pendimethalin

**Signal word:** CAUTION

**Formulation:**

Microencapsulated, suspension

**Use rates:** 3.1 - 6.3 pints per acre

**PPE:** Long-sleeved shirt, long pants, chemical-resistant gloves, shoes plus socks

**REI:** 24 hours

**Packaging:** 2.5 gal jug, 1 x 15 gal keg, 110 gal and 220 gal bulk totes



# Pendulum® 3.3 EC

## Herbicide

Nursery growers count on **Pendulum 3.3 EC** herbicide to deliver highly effective and economical preemergent control of the most challenging grassy and broadleaf weeds found in outdoor ornamental production.

### Key Features and Benefits

- May be used as a broadcast or directed spray, bare ground or over-the-top applications
- Extensively tested to provide excellent crop safety
- Controls more than 40 of the most challenging grassy and broadleaf weeds
- May be tank-mixed with preemergent **Tower®** herbicide for control of a broader range of weeds

### Weeds Controlled

Refer to the *Weeds Controlled* chart in this guide for common weeds controlled.

For a complete listing, visit [www.CDMS.net](http://www.CDMS.net) or find the product label using the convenient QR Code link below.

### Use Sites

- Field-, liner-, and container-grown ornamental production
- Bare ground areas for container placement
- Bulb plantings
- Non-bearing fruit and nut tree nursery production
- Conifer and hardwood seedling nursery production
- Tree plantations for site preparation and maintenance
- Residential and commercial landscape
- Golf courses
- Recreational parks and athletic fields
- Sod farms
- General grounds maintenance (non-cropland areas)



**WSSA group:** 3

**Active ingredient:**

Pendimethalin

**Signal word:** CAUTION

**Formulation:** EC

(Emulsifiable concentrate)

**Use rates:** 2.4-4.8 qt per acre

**PPE:** Coverall, chemical-resistant gloves, shoes plus socks

**REI:** 24 hours

**Packaging:** 2.5 gal jug, 1 x 15 gal keg





# Pendulum® 2G

Herbicide

**Pendulum 2G** is for the budget-conscious nursery grower and landscape manager who wants effective preemergent weed control in a spreadable granular formulation.

## Key Features and Benefits

- Controls more than 40 grassy and broadleaf weeds
- Proven safety testing over more than 335 ornamental species in outdoor nursery production and landscape settings
- Specifically formulated for consistent spreadability and distribution

## Weeds Controlled

Refer to the *Weeds Controlled* chart in this guide for common weeds controlled.

For a complete listing, visit [www.CDMS.net](http://www.CDMS.net) or find the product label using the convenient QR Code link below.

## Use Sites

- Field-, liner-, and container-grown ornamental production
- Bare ground areas for container placement
- Bulb plantings
- Non-bearing fruit and nut tree nursery production
- Conifer and hardwood seedling nursery production
- Tree plantations for site preparation and maintenance
- Residential and commercial landscape
- Golf courses
- Recreational parks and athletic fields
- Sod farms
- General grounds maintenance (non-cropland areas)

**WSSA group:** 3

**Active ingredient:**

Pendimethalin

**Signal word:** CAUTION

**Formulation:** G (Granular)

**Use rates:** 100-200 lb per acre in ornamentals

**PPE:** Coveralls, chemical-resistant gloves, shoes plus socks

**REI:** 24 hours

**Packaging:** 40 lb bag



# Tower®

## Herbicide

**Tower** preemergent liquid herbicide hits the mark for nursery growers who want to optimize their investment in spray applications and modern equipment technology for control of the toughest summer weeds.

### Key Features and Benefits

- Controls annual grasses, sedges and small-seeded broadleaf weeds
- Unsurpassed performance on difficult to control weeds such as spurge, groundsel, yellow nutsedge, kyllinga, doveweed
- More than 50 of the most challenging weeds listed on the label
- Excellent tank mix partner with **Pendulum® 3.3EC** or **Pendulum® AquaCap** to broaden the range of weeds controlled

### Weeds Controlled

Refer to the *Weeds Controlled* chart in this guide for common weeds controlled. For a complete listing, visit [www.CDMS.net](http://www.CDMS.net) or find the product label using the convenient QR Code link below.

### Use Sites

#### Commercial ornamental production

- Container nurseries
- Field grown nurseries
- Conifer, Christmas tree and hardwood liner nurseries and tree plantations
- Residential and commercial lawn and landscapes
- Golf courses
- Turfgrass



**WSSA group:** 15

**Active ingredient:**

Dimethenamid-P

**Signal word:** WARNING

**Formulation:** EC

(emulsifiable concentrate)

**Use rates:**

21-32 fl oz per acre

**PPE:** Coveralls, waterproof gloves, shoes plus socks, protective eyewear

**REI:** 12 hours

**Packaging:**

2 x 2.5 gal per case







# Finale®

## Herbicide

**Finale** herbicide gives you fast, effective control of a broad spectrum of emerged annual and perennial grasses, broadleaf weeds and woody species.

### Key Features and Benefits

- Kills weeds fast with visible results in 2-4 days under many conditions
- Controls glyphosate-resistant weeds
- Rainfast in 4 hours

### Weeds Controlled

Marestail, parthenium, pigweed, lambsquarter, ragweed, kochia, thistle, dandelion and poison ivy

For a complete listing, visit [www.CDMS.net](http://www.CDMS.net) or find the product label using the convenient QR Code link below.

### Use Sites

- In and around greenhouses
- Container nurseries
- Field nurseries
- Landscapes
- Bareground/non-crop areas
- Around Christmas trees

**WSSA group:** 10 =  
Inhibitor of glutamine  
synthetase

**Active ingredient:**  
Glufosinate-ammonium

**Signal word:** WARNING

**Formulation:** L (Liquid)

**Use rates:** 2-6 qts per  
acre

**PPE:** Coveralls, chemical-  
resistant gloves, shoes +  
socks, protective eyewear

**REI:** 12 hours

**Packaging:** 2 x 2.5 gallon

# Basagran® T&O

Herbicide

Ornamental growers who want to keep non-production and other open areas free of broadleaf weeds and annual sedges confidently choose

**Basagran T&O** postemergent herbicide for reliable, cost-effective control of emerged weeds.

## Key Features and Benefits

- Effective control of problematic weeds such as yellow nutsedge and kyllinga
- Controls thistles, wild mustard, lawn burweed and other difficult-to-control broadleaf weeds
- Tank-mix with **Tower®** herbicide to achieve preemergent control of a broader range of weeds
- May also be used in turf and landscape settings

## Weeds Controlled

Refer to the *Weeds Controlled* chart in this guide for common weeds controlled. For a complete listing, visit [www.CDMS.net](http://www.CDMS.net) or find the product label using the convenient QR Code link below.

## Use Sites

- Established turfgrass
- Ornamentals
- Nurseries
- Non-cropland sites, roadsides and rights-of-way



**WSSA group:** 6

**Active ingredient:**

Sodium bentazon

**Signal word:** CAUTION

**Formulation:** L (Liquid)

**Use rates:** 24-32 fl oz per acre

**PPE:** Coveralls, chemical-resistant gloves, shoes plus socks

**REI:** 48 hours

**Packaging:** 1 gal jug  
2 x 1.0 gal per case





# Segment® II

Herbicide

## Key Features and Benefits

- (safe) for all broadleaf plant species

## Weeds Controlled

Crabgrass, foxtail, johnsongrass, field sandbur, shattercane

For a complete listing, visit [www.CDMS.net](http://www.CDMS.net) or find the product label using the convenient QR Code link below.

## Use Sites

- Christmas tree and conifer nurseries and plantations
- Commercial and residential landscapes
- Container and field-grown nursery production
- Industrial landscapes
- Perennial peanuts (non food)
- Potting soil and top soil
- Forestry, reforestation and tree nurseries and plantations
- Turfgrass

**WSSA group:** 1

**Active ingredient:**

Sethoxydim

**Signal word:** WARNING

**Formulation:** EC

(Emulsifiable Concentrate)

**Use rates:** 1.5 - 2.5 pints per acre

**PPE:** Coveralls over short sleeve shirt and short pants, chemical-resistant gloves, chemical-resistant footwear + socks, protective eyewear, chemical-resistant headgear for overhead exposure

**REI:** 12 hours

**Packaging:**

## Herbicide Resistance Management Guide – Preemergent Herbicides

WSSA Group	Chemical Group(s)	Primary Target Weed(s)	Active Ingredient(s)	Herbicide Trade Name(s)	REI	Use Site(s)
3	<b>Benzamide</b>	Selected annual and broadleaf weeds	Pronamide	Kerb	24	N
	<b>Dinitroaniline</b>	Selected annual and broadleaf weeds	Pendimethalin	<b>Pendulum® 2G</b> herbicide, <b>Pendulum Aquacap</b> herbicide, <b>Pendulum 3.3 EC</b> herbicide, Pre-M®, ProPendi®, Corral® 2.68G	24	N,L
			Prodiamine	Barricade®, Regalkade®, Stonewall®	12	N,L
			Oryzalin	Surflan®, Weed Impede®	12	N,L
			Trifluralin	Treflan®	12	N,L
<b>Pyridine</b>		Dithiopyr	Dimension®, Lifeguard®	12	N,L	
12	<b>Pyridazinorle</b>	Selected annual & broadleaf weeds	Norflurazon	Predict	12	N
14	<b>N-Phenylphthalimide</b>	Selected annual grasses and broadleaf weeds	Flumioxazin	BroadStar® SureGuard®	12	N
	<b>Oxadiazole</b>	Selected annual grasses and broadleaf weeds	Oxadiazon	Ronstar®	12	N,L
15	<b>Acetamide</b>	Selected annual and broadleaf weeds	Napropamide	Devrinol®, Strawberry & Fruit Tree Weeder®	12	N,L
	<b>Chloroacetamide</b>	Yellow nutsedge and selected annual and broadleaf weeds	S-metolachlor	Pennant Magnum®	24	N,L
		Selected annual grasses, broadleaf weeds and annual sedges	Dimethenamid-P	<b>Tower®</b> herbicide	12	N,L
20	<b>Nitrile</b>	Selected annual and perennial weeds	Dichlobenil	Casoron®, Barrier®	12	N,L
21	<b>Benzamide</b>	Selected broadleaf weeds	Isoxaben	Gallery®	12	N,L
29	<b>Alkylazine</b>	Grasses, Annual sedge, Annual broadleaf weeds	Indaziflam	Marengo®, Marengo® G	12	G,N,L
3 + 3	<b>Dinitroaniline</b>	Selected annual and broadleaf weeds	Benefin + Oryzalin	XL 2G®	24	N,L
3 + 14	<b>Dinitroaniline + Oxadiazole</b>	Selected annual grasses and broadleaf weeds	Pendimethalin + Oxadiazon	Jewel®	12	N,L
3 + 15	<b>Dinitroaniline + Chloroacetamide</b>	Selected annual grasses, broadleaf weeds and annual sedges	Pendimethalin + Dimethenamid-P	<b>FreeHand® 1.75G</b> herbicide	12	N,L
21 + 3	<b>Benzamide + Dinitroaniline</b>	Selected annual and broadleaf weeds	Isoxaben + Trifluralin	Snapshot TG®, Gallery® Woodace Preen Plus®	12	N,L
			Isoxaben + Prodiamine	Gemini®, Gemini® G	12	N,L
	<b>Benzamide + Pyridine</b>	Selected annual and broadleaf weeds	Isoxaben + Dithiopyr	Fortress(R)	12	N
14 + 3	<b>Diphenylether + Dinitroaniline Oxadiazole + Dinitroaniline</b>	Selected annual and broadleaf weeds	Oxyfluorfen + pendimethalin	OH2® Rout®	24	N
			Oxyfluorfen + oryzalin	RegalStar®	12	N
			Oxyfluorfen + prodiamine	Biathalon®	24	N,L
			Oxyfluorfen + Oxadiazon	Regal 00®	24	N,L
21 + 14 + 3	<b>Benzamide + Oxadiazole + Dinitroaniline</b>	Selected annual and broadleaf weeds	Isoxaben + Oxyfluorfen + Trifluralin	Showcase®	24	N

G = Greenhouse, N= Nursery, L=Landscape



## Herbicide Resistance Management Guide – Postemergent Herbicides

WSSA Group	Chemical Group(s)	Primary Target Weed(s)	Active Ingredient(s)	Herbicide Trade Name(s)	REI	Use Site(s)
1	<b>Aryloxyphenoxy Propionate</b>	Annual and perennial grasses	Fluazifop-P	Fusilade® II, Ornamec®	12	N,L
			Fenoxaprop-P	Acclaim® Extra	12	N,L
	<b>Cyclohexanedione</b>		Sethoxydim	<b>Segment® II</b> herbicide	12	N,L
			Clethodim	Envoy® Envoy® Plus	12 24	G,N,L
2	<b>Sulfonylurea</b>	Annual and perennial sedges	Halosulfuron-methyl	Sedge Hammer®	12	L
4	<b>Pyridine carboxylic acid</b>	Broadleaf weeds, including Canada thistle, Legumes, Aster and Nightshade family	Clopyralid	Lontrel®	12	N,L
5	<b>Triazine</b>	Annual grassy and broadleaf weeds	Simazine	Princep®	12	N,L
6	<b>Benzothiadiazinone</b>	Yellow nutsedge, annual sedges and selected broadleaf weeds	Bentazon	<b>Basagran® T&amp;O</b> herbicide	48	N,L
9	<b>Glycine</b>	Annual and perennial weeds	Glyphosate	Gly-Flo®, GlyphoMate®, Gly Star® Pro, Prosecutor®, Roundup® Pro, Touchdown® Pro, Trail Blazer®	4	G,N,L
				Refuge	12	
10	<b>Phosphinic Acid</b>	Annual and perennial weeds	Glufosinate-ammonium	<b>Finale®</b> herbicide	12	N,L
14	<b>Triazolinone</b>	Selected annual and perennial broadleaf weeds	Carfentrazone-ethyl	Quicksilver®	NOL	L
			Sulfentrazone	Dismiss®	12	N,L
	<b>N-phenylphthalimide</b>		Flumioxazin	SureGuard®	12	G,N,L
20	<b>Nitrile</b>	Selected annual and perennial weeds	Dichlobenil	Casoron®, Barrier®	12	N,L
22	<b>Bipyridylum</b>	Annual and perennial weeds	Diquat	Reward®, Tribune®	24	G,N,L
			Paraquat	Gramoxone SL	12 - 24	
26	<b>Carboxylic acid</b>	Annual and perennial weeds	Pelargonic acid	Scythe®	24	G,N,L
9 + 22	<b>Glycine + Carboxylic acid</b>	Annual and perennial weeds	Glyphosate + Diquat	QuickPro® Roundup QuikPro®	NOL	L
NC	<b>Soaps, Fatty Acids</b>	Selected annual and perennial weeds	Ammoniated soaps of fatty acids	Quik-Fire®	24	G,N,L

G = Greenhouse, N= Nursery, L=Landscape

1 Some products within the same chemical group may not control all weeds listed, or there may be some products within the group that offer superior performance. Consult your local university/extension specialist if needed.

2 Specific use sites may vary by formulation and manufacturer; therefore, always read and consult product label directions.

**WSSA Classification List:**

- 1 Inhibition of acetyl CoA carboxylase (ACCase)
- 2 Inhibition of acetolactate synthase (ALS) or acetohydroxyacid synthase (AHAS)
- 3 Inhibition of microtubule assembly
- 4 Synthetic auxin
- 5 Inhibition of photosynthesis at photosystem II (site A)
- 6 Inhibition of photosynthesis at photosystem II (site B)
- 9 Inhibition of enolpyruvyl shikimate-3-phosphate (EPSP) synthase
- 10 Inhibition of glutamine synthetase
- 12 Inhibitor of phytoene desaturase (PDS)
- 14 Inhibition of protoporphyrinogen oxidase (Protox, PPO)
- 15 Inhibition of mitosis via inhibition of very long chain fatty acids (VLCFAs)
- 20 Inhibition of cell wall synthesis (site A)
- 21 Inhibition of cell wall synthesis (site B)
- 22 Inhibition of photosystem I
- 26 Unknown site of action
- 29 Inhibition of cell wall synthesis (site C)
- NC Not classified

*The above list was developed by the Weed Science Society of America (WSSA). WSSA is a non-profit professional society made up of scientists from academia and private industry that promote research, education and outreach in the field of Weed Science. More information on the Weed Science Society of America is available from [www.wssa.net](http://www.wssa.net).*











# Total Release Fungicides and Insecticides

Total release is a unique pesticide application system that delivers ULV (Ultra Low Volume) insecticides and fungicides throughout a greenhouse for effective pest control. Its exclusive aerosol delivery system reduces pesticide applicator exposure through a three-step process: *Locate*, *Lock Down*, and *Leave*. Total release products are ready to use, so you never have to mix the chemicals or operate industrial application equipment, and they effectively control a wide variety of insects and diseases that are common issues in the greenhouse. Total release aerosol delivery systems make pesticide applications simple and easy.

## Total Release Best Practices

How to Use the *Locate*, *Lock Down* and *Leave* application system:

### Step 1: Locate

Determine appropriate areas to place the containers.

- They work best on benches or hanging from trusses.
- Do not place containers directly in front of fans, heaters or inlets to jet tubes.
- After finding the best location, create a plant-free zone by removing any plants within the recommended clearance area, up to 6 feet around the container.
- Elevate the container so the top is above the height of the plants you will treat.
  - Place the container on an inverted empty pot at the proper elevation on the benches.
  - Ensure a plant-free, obstacle-free clearance zone around each container.
- If using multiple containers, plan your application placements and greenhouse exit path in advance to avoid walking through the pesticide fog.

### **Step 2: Lock Down**

Keep the area contained to create the best application environment for total release products.

- Close all vents and sources of outside air.
- Run horizontal air flow (HAF) fans or fan jets (no heat; air only) for at least 15 minutes prior to application to create air movement in the greenhouse.
- Shut down all HAF fans just before activating cans; HAF fans should remain off for at least three hours or, if possible, all night.
- Put on the required personal protection equipment (PPE) and hold the container at arm's length; position the tip so that the spray releases from the can at an angle away from the applicator.
- Depress the actuator until it locks down, and set the container back into the plant-free zone.
- If you are using multiple containers in the greenhouse, start at the farthest point and work your way toward the exit door, locking down containers as you proceed.

### **Step 3: Leave**

Once all the containers are locked down, leave the greenhouse and shut the exit door.

- The containers will fog for approximately 2 to 3 minutes until empty.
- Allow the pesticide particles to circulate for at least 4 hours.
- Common practice is to leave the house closed overnight; if necessary, HAF fans may be returned to operation after the minimum 4-hour circulation period. Refer to label for REI guidelines and ventilation requirements. ALWAYS FOLLOW LABEL DIRECTIONS.
- Once the REI is complete, you may enter the greenhouse and resume normal operations.
- If you need to re-enter the greenhouse before the REI, you must wear PPE as required by the label.

# Total Release Products

## Total Release Insecticides

### 1300 Orthene® TR

Micro total release insecticide

Ultra-low volume fogger, broad range insecticide for use in commercial greenhouses. Use on bedding plants, cut flowers, flowering hanging baskets, foliage, potted flowering plants and ornamentals.

**Key Insects Controlled:** Spider mites, thrips, whiteflies, aphids, mealybugs, fungus gnats, scale.

*For a complete list of insects controlled and product details, consult the product label at CDMS.net using the convenient QR Code link at right.*



Active Ingredient	IRAC Group	Mode of Action	Signal Word	REI	PPE	Packaging
12% acephate	Group 1A,1B <i>Organophosphate</i>	Acetylcholinesterase inhibitor	CAUTION	24 hours	See label	12 x 6 oz aerosol cans

### Attain® TR

Micro total release insecticide

Ultra-low volume fogger, broad range insecticide for use in commercial greenhouses. Use on bedding plants, cut flowers, flowering hanging baskets, foliage, potted flowering plants and ornamentals.

**Key Insects Controlled:** Mites, thrips, whiteflies, aphids, mealybugs, fungus gnats, scale.

*For a complete list of insects controlled and product details, consult the product label at CDMS.net using the convenient QR Code link at right.*



Active Ingredient	IRAC Group	Mode of Action	Signal Word	REI	PPE	Packaging
4% bifenthrin	Group 3 <i>Pyrethroid</i>	Sodium channel modulator	CAUTION	2-24 hours See label	See label	12 x 2 oz aerosol cans

### Beethoven® TR

Miticide/insecticide

Ultra-low volume fogger, insecticide for use in commercial greenhouses. Use on bedding plants, cut flowers, flowering hanging baskets, foliage, potted flowering plants and ornamentals. An excellent space mix partner with **Pylon TR** for quick knockdown, added insect pest control and resistance management.

**Key Insects Controlled:** Spider mites, suppression of whiteflies.

*For a complete list of insects controlled and product details, consult the product label at CDMS.net using the convenient QR Code link at right.*



Active Ingredient	IRAC Group	Mode of Action	Signal Word	REI	PPE	Packaging
5% etoxazole	Group 10	Mite growth inhibitor	CAUTION	2-24 hours See label	See label	12 x 2 oz aerosol cans

### Duraplex® TR

Micro total release insecticide

RESTRICTED USE PESTICIDE. Refer to label for specific use conditions. Ultra-low volume fogger, broad range insecticide for use in commercial greenhouses. Use on bedding plants, cut flowers, flowering hanging baskets, foliage, potted flowering plants and ornamentals.

**Key Insects Controlled:** Mites, thrips, whiteflies, aphids, mealybugs, fungus gnats, shore flies, scale.

*For specific use conditions, a complete list of insects controlled and product details, consult the product label at CDMS.net using the convenient QR Code link at right.*



Active Ingredient	IRAC Group	Mode of Action	Signal Word	REI	PPE	Packaging
8% chlorpyrifos 1.6% cyfluthrin	Group 3A, 1B	Sodium channel modulator, Acetylcholinesterase inhibitor	WARNING	24 hours	See label	12 x 2 oz aerosol cans

## Total Release Products *(continued)*

### Total Release Insecticides *(continued)*

#### **Pylon® TR**

Total release insecticide

Ultra-low volume fogger insecticide for use in commercial greenhouses and fully enclosed garden centers. Use on bedding plants, cut flowers, flowering hanging baskets, foliage, potted flowering plants, fruiting vegetables, and ornamentals. An excellent space mix partner with **Beethoven® TR** for quick knockdown, added insect pest control and resistance management.

**Key Insects Controlled:** Mites, spider mites, thrips, fungus gnats.

*For a complete list of insects controlled and product details, consult the product label at [CDMS.net](http://CDMS.net) using the convenient QR Code link at right.*



Active Ingredient	IRAC Group	Mode of Action	Signal Word	REI	PPE	Packaging
4.5% chlorfenapyr	Group 13 Pyrrole	Interrupts energy production in cell mitochondria	WARNING	2-12 hours See label	See label	12 x 2 oz aerosol cans

#### **Pyrethrum® TR**

Micro total release insecticide

Ultra-low volume fogger, broad range insecticide for use in commercial greenhouses. Use on bedding plants, cut flowers, flowering hanging baskets, foliage, potted flowering plants, specified vegetables (root and leafy), fruiting vegetables, herbs, and ornamentals. An excellent space mix partner with **1300 Orthene® TR** or **Duraplex® TR** for added insect pest control and resistance management. When used with another Total Release insecticide, the most restrictive label applies. DO NOT combine with another Total Release product when applied to vegetables and herbs.

**Key Insects Controlled:** Mites, thrips, whiteflies, aphids, mealybugs, fungus gnats, shore flies, scale.

*For a complete list of insects controlled and product details, consult the product label at [CDMS.net](http://CDMS.net) using the convenient QR Code link at right.*



Active Ingredient	IRAC Group	Mode of Action	Signal Word	REI	PPE	Packaging
4% pyrethrin 16% piperonyl butoxide	Group 3a Pyrethroid	Botanical, sodium channel modulator	Caution	4-12 hours	See label	12 x 2 oz aerosol cans

## Total Release Products *(continued)*

### Total Release Fungicides

**Pageant® TR**  
Total release  
fungicide

Ultra-low volume fogger for broad spectrum disease control in commercial greenhouses and transportation equipment (trucks-cargo areas only). Use on bedding plants, cut flowers, flowering hanging baskets, foliage plants, potted flowering plants and perennials.



**Diseases Controlled:** *Alternaria* spp., *Botrytis* spp., *Cercospora*, *Cylindrocladium* spp., *Myrothecium* spp., Powdery Mildew, and Rust (*Uromyces*)

For a complete list of diseases controlled and product details, consult the product label at [CDMS.net](http://CDMS.net) using the convenient QR Code link at right.

Active Ingredient	FRAC Group	Mode of Action	Signal Word	REI	PPE	Packaging
3.50% Boscalid	7	Succinate-dehydrogenase enzyme inhibitor (SDHI)	Caution	4-12 hours	See Label	12 x 3 oz aerosol can
1.75% Pyraclostrobin	11	Quinone outside Inhibitor (QoI)				

**Trinity® TR**  
Total release  
fungicide

Ultra-low volume fogger, broad-spectrum foliar disease control for use in commercial greenhouses. Use on bedding plants, cut flowers, flowering hanging baskets, foliage, potted flowering plants, perennials and other ornamentals.



**Diseases Controlled:** *Alternaria* spp., *Botrytis* spp., *Cercospora*, *Cylindrocladium* spp., *Fusarium* spp., *Myrothecium* spp., Powdery mildew and *Rhizoctonia* spp.

For a complete list of diseases controlled and product details, consult the product label at [CDMS.net](http://CDMS.net) using the convenient QR Code link at right.

Active Ingredient	FRAC Group	Mode of Action	Signal Word	REI	PPE	Packaging
10.5% triticonazole	Group 3 Triazoles	Demethylation inhibitor, (DMI)	CAUTION	4-12 hours See label	See label	12 x 3 oz aerosol cans









# Specialty Greenhouse Products

Rely on the BASF portfolio of specialty products to address a variety of production needs in greenhouse operations, from identifying and monitoring pest populations to safely sanitizing equipment and facilities. These products offer additional layers of pest, disease and application control to help improve quality, plant health and product performance.

## Specialty Greenhouse Product Use Site Recommendations

Product	Bareground	Conifer Nursery Plantation	Container Nursery	Field Production	Retail Nursery	Greenhouse	Interior	Landscape and Recreational Areas	Turf
<b>Pest Control</b>									
<b>PT® Wasp Freeze®</b> Wasp & Hornet Insecticide	■	■	■	■	■	■	■	■	■
<b>Iron Chelates</b>									
<b>Sprint® 138</b> Iron Chelates		■	■	■	■	■	■	■	■
<b>Sprint® 330</b> Iron Chelates		■	■	■	■	■	■	■	■
<b>Disinfectant and Algicide</b>									
<b>Green-Shield® II</b> Disinfectant and Algicide						■	■ <sup>1</sup>		
<b>Insect Monitoring</b>									
<b>Sensor®</b> Insect Monitoring and Trapping Cards		■	■	■	■	■	■		
<b>Spray Pattern Indicators</b>									
<b>Turf Mark®</b> Spray Pattern Indicator	■	■	■	■		■		■	■
<b>Tank Cleaners</b>									
<b>Nutra-Sol®</b> Tank Cleaner	■	■	■	■	■	■	■	■	■
<b>Neutralize®</b> Tank Cleaner	■	■	■	■	■	■	■	■	■
<b>Other</b>									
<b>Instant Soap 'N Water®</b> Foaming Cleanser <sup>2</sup>	■	■	■	■	■	■	■	■	■

*covers to be removed on a seasonal basis.*

<sup>2</sup> Consult Green-Shield II label for interior uses.

<sup>3</sup> Instant Soap 'N Water foaming cleanser is labeled for use on hands in all use sites listed.





# Sprint® 138

## Iron Chelate

A strongly chelated iron micronutrient that produces superior results regardless of soil pH, **Sprint 138** iron chelates help horticultural professionals solve plant health issues related to iron chlorosis.

### Key Features and Benefits

- Prevents and corrects iron deficiency
- Strongest chelate available for the most challenging alkaline or calcareous soils
- Works across a wide range of ornamental plant varieties in horticulture production settings
- Formulated to maintain and protect iron availability across all pH soils
- Flexible use as either foliar or soil applications

### Keys to Success

- Optimize efficacy with good foliar spray coverage or directed placements at the root zone, or dry incorporation into soil media
- Follow with irrigation or time applications to precede rainfall events
- Ensure thorough mixing and compatibility in spray solution combinations with other crop protection products and fertilizers

### Use Sites

- Production greenhouses
- Nursery horticulture
- Professional turf management
- Commercial landscapes
- Residential landscapes
- Flower beds
- Shrubs
- Fruit trees
- Ornamental trees
- Vegetables

#### Active ingredient:

Chelated Iron

#### Formulation:

Dry powder

#### Use rates:

*Ornamentals:* Refer to product package label for specific applications to ornamental plants, trees, shrubs, vegetables, and/or application table for mechanical injection rates

*Turf:* 2 - 4 oz per 1000 sq ft in sufficient water to achieve uniform coverage

**Packaging:** 6 x 5 lb bags





# Sprint® 330

## Iron Chelate

**Sprint 330** iron chelate corrects iron micronutrient deficiency in plants to achieve better color and plant quality. Unlike non-chelated or weakly-chelated iron products, **Sprint 330** iron chelate provides for more efficient plant uptake, producing long-lasting vibrant color.

### Key Features and Benefits

- Maintains and protects iron availability in slightly acidic to slightly alkaline soils
- Long-proven professional performance in production horticulture
- Corrects iron deficiency across a wide range of ornamental varieties
- Flexible, economical use as either foliar or soil applications

### Keys to Success

- Performs best in soils where pH is 6.0 - 7.5
- Optimize efficacy with good foliar spray coverage or directed placements at the root zone, or dry incorporation into soil media
- Follow with irrigation or time applications to precede rainfall
- Ensure thorough mixing and compatibility in spray solution combinations with other crop protection products and fertilizers – if necessary, conduct jar test

### Use Sites

- |                                |                    |
|--------------------------------|--------------------|
| ■ Production greenhouses       | ■ Flower beds      |
| ■ Nursery horticulture         | ■ Shrubs           |
| ■ Professional turf management | ■ Fruit trees      |
| ■ Commercial landscapes        | ■ Ornamental trees |
| ■ Residential landscapes       | ■ Vegetables       |



### Active ingredient:

Chelated iron

### Formulation:

Dry powder

### Use rates:

*Ornamentals:* Refer to product package label for specific applications to ornamental plants, trees, shrubs, vegetables, and/or application table for mechanical injection rates

*Turf:* 1 - 4 oz per 1000 sq ft in sufficient water to achieve uniform coverage

**Packaging:** 6 x 5 lb bags, 1 x 50 lb box





# Green-Shield® II

## Disinfectant and Algicide

Recognized for its consistently high quality, **Green-Shield II** disinfectant and algicide provides growers with a biodegradable, cost-effective treatment for the control of unwanted pathogens and algae in greenhouse operations.

**Group:**

Quaternary ammonium

**Active ingredient:**

Quaternary amines

**Signal word:** DANGER

**Formulation:**

Liquid

**Use rates:** Various—refer to product label for specific treatments and applications.

**PPE:** Coveralls worn over long-sleeved shirt and long pants; waterproof gloves; chemical-resistant footwear plus socks; protective eyewear (goggles or face shield); chemical-resistant headgear. When using this product for algae control in overhead watering systems, wear a chemical-resistant apron when mixing, loading or cleaning equipment.

**REI:** None

**Packaging:** 4 x 1 gal, 30 gal drums

**Key Features and Benefits**

- Effectively disinfects and sanitizes against bacteria and fungus on walkways, greenhouse coverings and evaporative cooling pads
- Broad-spectrum control of pathogens and algae found on floors, benches and walls in and around greenhouses
- Less irritating to use with no odor, unlike bleach solutions

**Keys to Success**

- Use to disinfect work areas, hand tools and containers to prevent transfer of pathogens
- Use every other week to control algae and odor in evaporative cooling pools
- Use during the cropping cycle for routine disinfection and algae control on benches and walkways
- Use in production areas in cleanup program during transition periods

**Use Sites**

- Greenhouses
- Nurseries
- Landscapes and interiorscapes on work areas
- Benches
- Tools
- Containers
- Walkways
- Floors
- Evaporative cooling pools and pads



# Sensor®

## Insect Monitoring and Trapping Cards

**Sensor** insect monitoring and trapping cards are an important component of effective IPM strategies to help growers achieve effective pest population monitoring and identification in production operations.

### Key Features and Benefits

- Use to monitor changing insect populations
- Easy-to-remove covers
- Gridded for easy reading
- Durable, long-lasting

### Keys to Success

- Begin with positioning one card per 500 sq ft
- Use additional cards to monitor entry points such as doors and vents
- Monitor “hot spots” within susceptible ornamental varieties
- Record insect catch data on a routine basis during the growing season
- Replace cards at regular intervals and always after spraying insecticides

### Use Sites

- Greenhouses
- Nurseries
- Laboratories




### Formulation:

Glue card

**Use rates:** Start with one card per 500 sq ft. Use additional cards to monitor entry points (doors and vents), to determine “hot” spots within susceptible varieties, and to monitor low-level populations. Change cards at routine intervals and always after spraying insecticides. Record insect catch data on a routine basis.

**Packaging:** 750 cards per case (50 cards per pack, 15 packs per case)



# Turf Mark®

## Spray Pattern Indicator

**Turf Mark** brand colorants are used by professional spray applicators who want to make visible and precise pesticide and fertilizer applications in production horticulture, turf management, and landscape maintenance.

### Key Features and Benefits

- Helps prevent application skips and overlaps
- Shows application pattern for increased accuracy and coverage, improving service quality and product performance
- Colorant biodegrades quickly under sunlight and moisture

### Keys to Success

- Use in horticulture production stock tank solutions as a color indicator in irrigation water delivered to plant containers
- Choose Turf Mark Green colorant to achieve more discrete applications in turf and landscape
- Avoid overspray onto building walls, decorative hardscapes, driveways, sidewalks and on other hard porous materials where residual discoloration is undesirable

### Use Sites

- Greenhouses
- Nurseries
- Residential landscapes
- Commercial landscapes
- Residential turf
- Commercial turf
- Bare ground
- Fence lines
- Roadsides

**Active ingredient:**

Dye Colorant

**Formulation:**

Liquid or dry powder

**Use rates:** 1-2 qt

per 100 gal

**Packaging:** 12 x 1 qt,

4 x 1 gal, 2 x 2.5 gal,

1 x 30 gal drums,

40 x1 WSP



# Nutra-Sol™

Tank Cleaner

# Neutralize™

Tank Cleaner

**Nutra-Sol** and **Neutralize** tank cleaners allow professional applicators to safely and effectively remove fertilizer and pesticide residues from spray equipment. Use with confidence to clean all working parts, including tanks, pumps, hoses, valves and nozzles.

## Key Features and Benefits

- Cleans and removes residues left by 2,4-D, sulfonylureas, and other hard-to-remove residues that are active at low rates
- Non-corrosive formulations are safe to use on spraying systems, inside and out
- Prevents damage to rubber, plastic and other soft sprayer parts

## Keys to Success

- An end-of-work-day thorough rinsing and cleaning helps maintain integrity of spray equipment
- Removal of aggressive chemical residues helps prevent damage to rubber, plastic, and other soft sprayer parts
- Practice “finish clean, start clean”

## Use Sites

All spray application equipment used in greenhouses, nurseries, residential and commercial turf and landscape maintenance



**Active ingredient:**  
Surfactants

**DHS Signal word:**  
DANGER

**Formulation:**  
Dry powder, liquid

**Use rates:**  
1 - 2 lb per 100 gals  
1 quart per 100 gals

**Precautionary State-ments (Prevention):** Wear protective gloves, protective clothing, eye protection, face protection. Do not breathe dust or mist.

Refer to product label for complete information.

**Packaging:** 12 x 2 lb jars, 4 x 1 gal







# PT Wasp-Freeze® II

## Wasp & Hornet Insecticide

For protection of outdoor workers who need powerful knockdown and control of stinging insects, **PT Wasp Freeze II** wasp and hornet insecticide is conveniently applied to reach nests in even the toughest locations.

### Key Features and Benefits

- Contact activity for instant knockdown
- Works from up to 15 feet away
- Can be used around electrical equipment
- Contains no ozone-depleting substances

### Key Insects Controlled

- Wasps
- Hornets
- Yellowjackets
- Bees
- Spiders

### Use Site

- Outdoors

**IRAC group:** 3A

**Active ingredients:**

Prallethrin

**Signal word:** CAUTION

**Formulation:**

Aerosol

**Use rates:**

Directed spray-to-wet for immediate knockdown, as needed

**PPE:** None required

**REI:** None

**Packaging:**

12 x 17.5 oz cans



*Revere is a registered trademark of ADAMA. Gly Star is a registered trademark of Albaugh, LLC/Agri Star. Ornazin is a trademark and Durham, Terraclor and Turfcide are registered trademarks of Amvac Chemical Corporation. Decree, Dimilin and Endorse are registered trademarks of Arysta LifeScience Canada. Casoron, Granuflo and Kaligreen are registered trademarks of Arysta LifeScience North America LLC. AquaCap, Better Plants, and Orkestra are trademarks and Basagran, Beethoven, Empress, FreeHand, Green-Shield, Insignia, Intrinsic, Nemasys, Mil-lenium, Orvego, Pageant, Pendulum, Pylon, Segment, Siesta, Sprint, Stature, Sultan, and Tower are registered trademarks of BASF. CoreTect, Savate, Signature, TopChoice and Trigo are trademarks, and 26GT, Aliette, Allectus, Banol, Bayleton, Chipco, DeltaGard, Dylox, FenStop, Finale, Forbid, Interface, Kontos, Merit, Mirage, Prostar, Specticle, Stressgard, Tartan, Tempo and Thiodan are registered trademarks of Bayer Environmental Science. TerraClean and ZeroTol are registered trademarks of BioSafe Systems LLC. BotaniGard, Plantshield, Cease, Milstop, SuffOil and TriCon are registered trademarks of BioWorks Inc. Amdro is a registered trademark of Central Garden and Pet Company. Deliver, Javelin, Sluggo and Soilgard is a registered trademark of Certis USA, LLC. Terrazole, Terraguard, Adept and Floramite are registered trademarks of Chemtura Corporation. Affirm, Alude, Protect and Torque are trademarks and 3336, Spectro (90) and TriStar are registered trademarks of Cleary Chemical LLC. Dimension, Dithane, Dursban, Entrust, Fore, Eagle, Gallery, Intrepid, Lontrel, Rainshield, Snapshot and Conserve are registered trademarks of Dow AgroSciences LLC. Kocide is a registered trademark of DuPont Crop Protection. EV-N-SPRED is a registered trademark of EarthWay Products, Inc. Banrot and Ovation are registered trademarks of Everris International BV. Aria, Cygon and Segway and are trademarks and Glyphos, Malathion, OnyxPro and Talstar are registered trademarks of FMC Corporation. Astro and Quicksilver are registered trademarks of FMC Professional Solutions. Showcase is a trademark and Captiva, Confirm, Hexygon, Imidan, M-Pede, MSR Spray Concentrate, Magus, Mesuroil, Rubigan, Sanmite, Scythe, SledgeHammer, Supracide, Treflan and Ultiflora are registered trademarks of Gowan Company. Pedestal is a registered trademark of Makhteshim Agan. Regalia is a registered trademark of Marrone Bio Innovations. Roundup is a registered trademark of Monsanto. Actinovate is a registered trademark of Novosymes BioAg Inc. Emblem and Strider are trademarks and 26/36 and Tourney are registered trademarks of NuFarm Specialty Products. Applause, FenStop, Decathlon, Judo, Shuttle and Sirocco are trademarks and Azatin, Compass, Discus, Fulcrum, Marathon, Marengo, Segway, Strike (Plus) and Triact are registered trademarks of OHP, Inc. Orthene and OH2 are registered trademarks of OMS Investment, Inc. Barrier, GlyphoMate and Ornamec are registered trademarks of PBI-Gordon Professional. PureSpray is a trademark of Petro-Canada. Camelot and Promite are trademarks and Pen-tathlon and Talus are registered trademarks of SePRO Corporation. Rout is a registered trademark of The Scotts Company. Fusilade, Mural and Princep are trademarks of Syngenta Crop Protection, LLC. Enfold, Flagship, Hurricane and Micora are trademarks and Advion, Award, Alamo, Avid, Banner MAXX, Barricade, Citation, Daconil Ultrex, Endeavor, Heritage, Mainspring, MAXX, Medallion, Pennant MAGNUM, Reward, Scimitar, Subdue, Segovis and Touchdown are registered trademarks of Syngenta Professional Products. Fungo, perm-Up and Surflan are registered trademarks of United Phosphorus, Inc. Aloft, BroadStar, Envoy, Overture and TetraSan are trademarks and Adorn, Safari, Distance, Gnatrol, DiPel, SureGuard, Tame and Tourney are registered trademarks of Valent U.S.A. Corporation. Enstar, Extinguish, Mavrik and Enstar are registered trademarks of Wellmark International. Attain, Duraguard, Duraplex, Green-Shield, Sensor, Wasp-Freeze and X-clude are registered trademarks of Whitmire-Microgen Research Laboratories, Inc. Corral is a registered trademark of Winfield Solutions LLC.*

THE DESCRIPTIONS, DESIGNS, DATA AND INFORMATION CONTAINED HEREIN ARE PRESENTED IN GOOD FAITH, AND ARE BASED ON BASF'S CURRENT KNOWLEDGE AND EXPERIENCE. THEY ARE PROVIDED FOR GUIDANCE ONLY, AND DO NOT CONSTITUTE THE AGREED CONTRACTUAL QUALITY OF THE PRODUCT OR A PART OF BASF'S TERMS AND CONDITIONS OF SALE. BECAUSE MANY FACTORS MAY AFFECT PROCESSING OR APPLICATION/USE OF THE PRODUCT, BASF RECOMMENDS THAT THE READER CARRY OUT ITS OWN INVESTIGATIONS AND TESTS TO DETERMINE THE SUITABILITY OF A PRODUCT FOR ITS PARTICULAR PURPOSE PRIOR TO USE. IT IS THE RESPONSIBILITY OF THE RECIPIENT OF PRODUCT TO ENSURE THAT ANY PROPRIETARY RIGHTS AND EXISTING LAWS AND LEGISLATION ARE OBSERVED. **NO WARRANTIES OF ANY KIND, EITHER EXPRESS OR IMPLIED, INCLUDING, BUT NOT LIMITED TO, WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, ARE MADE REGARDING PRODUCTS DESCRIBED OR DESIGNS, DATA OR INFORMATION SET FORTH HEREIN, OR THAT THE PRODUCTS, DESCRIPTIONS, DESIGNS, DATA OR INFORMATION MAY BE USED WITHOUT INFRINGING THE INTELLECTUAL PROPERTY RIGHTS OF OTHERS.** ANY DESCRIPTIONS, DESIGNS, DATA AND INFORMATION GIVEN IN THIS PUBLICATION MAY CHANGE WITHOUT PRIOR INFORMATION. THE DESCRIPTIONS, DESIGNS, DATA, AND INFORMATION FURNISHED BY BASF HEREUNDER ARE GIVEN GRATIS AND BASF ASSUMES NO OBLIGATION OR LIABILITY FOR THE DESCRIPTIONS, DESIGNS, DATA OR INFORMATION GIVEN OR RESULTS OBTAINED, ALL SUCH BEING GIVEN AND ACCEPTED AT THE READER'S RISK.



We create chemistry

**BASF**

26 Davis Drive  
Research Triangle Park, NC 27709  
USA  
[betterplants.basf.us](http://betterplants.basf.us)  
[@BASFPlants\\_us](https://twitter.com/BASFPlants_us)

Always read and follow label directions.

©2019 BASF Corporation.

All rights reserved.

PSS 19-30-449 r07/19